

- 1 -

5

INHIBITORS OF PHOSPHOLIPASE ENZYMES

This application is a continuation-in-part of co-pending U.S. Serial No. 09/256,413, filed February 24, 1999, which claims the benefit of U.S. Provisional Application No. 60/100,426, which was converted from U.S. Patent Application No. 10 09/030,062, filed February 28, 1998, pursuant to a petition filed under 37 C.F.R. 1.53(c)(2)(i).

Background of the Invention

15 The present invention relates to chemical inhibitors of the activity of various phospholipase enzymes, particularly phospholipase A₂ enzymes.

Leukotrienes and prostaglandins are important mediators of inflammation, each of which classes contributes to the development of an inflammatory response in a different way. Leukotrienes recruit inflammatory cells such as neutrophils to an 20 inflamed site, promote the extravasation of these cells and stimulate release of superoxide and proteases which damage the tissue. Leukotrienes also play a pathophysiological role in the hypersensitivity experienced by asthmatics [See, e.g. B. Samuelson et al., Science, 237:1171-76 (1987)]. Prostaglandins enhance 25 inflammation by increasing blood flow and therefore infiltration of leukocytes to inflamed sites. Prostaglandins also potentiate the pain response induced by stimuli.

Prostaglandins and leukotrienes are unstable and are not stored in cells, but are instead synthesized [W. L. Smith, Biochem. J., 259:315-324 (1989)] from 30 arachidonic acid in response to stimuli. Prostaglandins are produced from arachidonic acid by the action of COX-1 and COX- 2 enzymes. Arachidonic acid is

- 2 -

5 also the substrate for the distinct enzyme pathway leading to the production of leukotrienes.

Arachidonic acid which is fed into these two distinct inflammatory pathways is released from the sn-2 position of membrane phospholipids by phospholipase A₂ enzymes (hereinafter PLA₂). The reaction catalyzed by PLA₂ is believed to represent the rate-limiting step in the process of lipid mediated biosynthesis and the production of inflammatory prostaglandins and leukotrienes. When the phospholipid substrate of PLA₂ is of the phosphatidyl choline class with an ether linkage in the sn-1 position, the lysophospholipid produced is the immediate precursor of platelet activating factor (hereafter called PAF), another potent mediator of inflammation [S.I. Wasserman, Hospital Practice, 15:49-58 (1988)].

Most anti-inflammatory therapies have focussed on preventing production of either prostaglandins or leukotrienes from these distinct pathways, but not on all of them. For example, ibuprofen, aspirin, and indomethacin are all NSAIDs which inhibit the production of prostaglandins by COX-1/COX-2, but have no effect on the inflammatory production of leukotrienes from arachidonic acid in the other pathways. Conversely, zileuton inhibits only the pathway of conversion of arachidonic acid to leukotriene, without affecting the production of prostaglandins. None of these widely-used anti-inflammatory agents affects the production of PAF.

Consequently the direct inhibition of the activity of PLA₂ has been suggested as a useful mechanism for a therapeutic agent, i.e., to interfere with the inflammatory response. [See, e.g., J. Chang et al, Biochem. Pharmacol., 36:2429-2436 (1987)].

5 A family of PLA₂ enzymes characterized by the presence of a secretion signal sequenced and ultimately secreted from the cell have been sequenced and structurally defined. These secreted PLA₂s have an approximately 14 kD molecular weight and contain seven disulfide bonds which are necessary for activity. These PLA₂s are found in large quantities in mammalian pancreas, bee venom, and various snake
10 venom. [See, e.g., references 13-15 in Chang et al, cited above; and E. A. Dennis, Drug Devel. Res., 10:205-220 (1987).] However, the pancreatic enzyme is believed to serve a digestive function and, as such, should not be important in the production of the inflammatory mediators whose production must be tightly regulated.

15 The primary structure of the first human non-pancreatic PLA₂ has been determined. This non-pancreatic PLA₂ is found in platelets, synovial fluid, and spleen and is also a secreted enzyme. This enzyme is a member of the aforementioned family. [See, J. J. Seilhamer et al, J. Biol. Chem., 264:5335-5338 (1989); R. M. Kramer et al, J. Biol. Chem., 264:5768-5775 (1989); and A. Kando et al, Biochem.
20 Biophys. Res. Comm., 163:42-48 (1989)]. However, it is doubtful that this enzyme is important in the synthesis of prostaglandins, leukotrienes and PAF, since the non-pancreatic PLA₂ is an extracellular protein which would be difficult to regulate, and the next enzymes in the biosynthetic pathways for these compounds are intracellular proteins. Moreover, there is evidence that PLA₂ is regulated by protein kinase C and
25 G proteins [R. Burch and J. Axelrod, Proc. Natl. Acad. Sci. U.S.A., 84:6374-6378 (1989)] which are cytosolic proteins which must act on intracellular proteins. It would be impossible for the non-pancreatic PLA₂ to function in the cytosol, since the high reduction potential would reduce the disulfide bonds and inactivate the enzyme.

30 A murine PLA₂ has been identified in the murine macrophage cell line, designated RAW 264.7. A specific activity of 2 mols/min/mg, resistant to reducing

1003035 001202

- 4 -

5 conditions, was reported to be associated with the approximately 60 kD molecule. However, this protein was not purified to homogeneity. [See, C. C. Leslie et al, Biochem. Biophys. Acta., 963:476-492 (1988)]. The references cited above are incorporated by reference herein for information pertaining to the function of the phospholipase enzymes, particularly PLA₂.

10

A cytosolic phospholipase A₂ (hereinafter "cPLA₂") has also been identified and cloned. See, U.S. Patent Nos. 5,322,776 and 5,354,677, which are incorporated herein by reference as if fully set forth. The enzyme of these patents is an intracellular PLA₂ enzyme, purified from its natural source or otherwise produced in
15 purified form, which functions intracellularly to produce arachidonic acid in response to inflammatory stimuli.


Now that several phospholipase enzymes have been identified, it would be desirable to identify chemical inhibitors of the action of enzymes, which inhibitors
20 could be used to treat inflammatory conditions, particularly where inhibition of production of prostaglandins, leukotrienes and PAF are all desired results. There remains a need in the art for an identification of such anti-inflammatory agents for therapeutic use in a variety of disease states.

25

Summary of the Invention

Compounds of this invention have the following formulae:


- 5 -



wherein:

R_1 and $R_{1'}$ are independently selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_{10}$ alkyl, preferably $-C_1-C_6$ alkyl, $-S-C_1-C_{10}$ alkyl, preferably $-S-C_1-C_6$ alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-CN$, $-NO_2$, $-NH_2$, $-HN(C_1-C_6)$, $-N(C_1-C_6)_2$, phenyl, $-O$ -phenyl, $-S$ -phenyl, benzyl, $-O$ -benzyl, or $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$;

or R_1 and $R_{1'}$ are a moiety of the formulae:



- 6 -

5 R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $C(O)CH_3$, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$;

10 R_7 is selected from $-OH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-N(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, quinolyl, $(CH_2)_n$ phenyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, $-(CH_2)_n$ -phenyl-O-phenyl, $-(CH_2)_n$ -phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ -phenyl-
15 $(O-CH_2$ -phenyl) $_2$, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NH_2$, $-NO_2$, $-COOH$, $-CF_3$, CO_2H , or $-OH$;


n is an integer from 0 to 3;

20


R_2 is selected from H, halogen, $-CF_3$, $-OH$, $-C_1$ - C_{10} alkyl, preferably $-C_1$ - C_6 alkyl, C_1 - C_{10} alkoxy, preferably C_1 - C_6 alkoxy, $-CHO$, $-CN$, $-NO_2$, $-NH_2$, $-NH-C_1$ - C_6 alkyl, $-N(C_1-C_6 \text{ alkyl})_2$, $-N-SO_2-C_1-C_6$ alkyl, or $-SO_2-C_1-C_6$ alkyl;

25

R_3 is selected from H, $-CF_3$, $-COOH$, C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, C_3 - C_{10} cycloalkyl, $-C_1$ - C_6 alkyl- C_3 - C_{10} cycloalkyl, $-CHO$, halogen, or a moiety of the formulae:




- 7 -




10 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₁-C₆ alkyl, C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O,
 15 preferably S or O;

R₄ is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, -(CH₂)_n-C₃-C₆ cycloalkyl, -(CH₂)_n-S-(CH₂)_n-C₃-C₅ cycloalkyl, -(CH₂)_n-O-(CH₂)_n-C₃-C₅ cycloalkyl, or the groups of:

20 a) -(CH₂)_n-phenyl-O-phenyl, -(CH₂)_n-phenyl-CH₂-phenyl, -(CH₂)_n-O-phenyl-CH₂-phenyl, -(CH₂)_n-phenyl-(O-CH₂-phenyl)₂, or a moiety of the formulae:




- 8 -



wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O; or

b) a moiety of the formulae -(CH₂)_n-A, -(CH₂)_n-S-A, or -(CH₂)_n-O-A, wherein A is the moiety:




wherein

D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;

B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, or -NO₂; or

c) a moiety of the formulae:

- 9 -



10 wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$; or

d) a moiety of the formula $-\text{L}^2\text{-M}^2$, wherein:

- 10 -

5

L^2 indicates a linking or bridging group of the formulae $-(CH_2)_n-$, $-S-$, $-O-$, $-SO_2-$, $-C(O)-$, $-(CH_2)_n-C(O)-$, $-(CH_2)_n-C(O)-(CH_2)_n-$, $-(CH_2)_n-O-(CH_2)_n-$, or $-(CH_2)_n-S-(CH_2)_n-$, $-C(O)C(O)X$;

where $X = O, N$

10

M^2 is selected from the group of C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, or $-CF_3$; or

15

i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, or $-CF_3$; or

20

ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-CHO$, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$ or $-OH$; or

25



30

- 11 -


- 5 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;
- 10

n is an integer from 0 to 3;

- R₅ is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, (CH₂)_n-CH=CH-COOH, -(CH₂)_n-tetrazole, -CH₂-phenyl-C(O)-benzothiazole,
- 15



, or




- 20 or a moiety selected from the formulae -L¹-M¹;

wherein L¹ is a bridging or linking moiety selected from a chemical bond, -(CH₂)_n-, -S-, -O-, -SO₂-, -C(O)-, -(CH₂)_n-C(O)-, -(CH₂)_n-C(O)-(CH₂)_n-, -(CH₂)_n-O-(CH₂)_n-, (CH₂)_n-S-(CH₂)_n-, -C(Z)-N(R₆)-, -C(Z)-N(R₆)-(CH₂)_n-, -C(O)-C(Z)-N(R₆)-, -C(O)-

- 12 -

- 5 $C(Z)-N(R_6)-(CH_2)_n-$, $-C(Z)-NH-SO_2-$, $-C(Z)-NH-SO_2-(CH_2)_n-$, $-(CH_2)_n-S-(CH_2)_n-$, $-(CH_2)_n-SO-(CH_2)_n-$, $-(CH_2)_n-SO_2-(CH_2)_n-$, or $-CH_2-CH=CH-CH_2-O-$;

M^1 is selected from the group of $-COOH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, tetrazole,




where R_8 , R_9 or R_{10} can be attached anywhere in the cyclic or bicyclic system,
 n is an integer from 0 to 3;

- 13 -

5

R_8 , in each appearance, is independently selected from H, -COOH, $-(CH_2)_n$ -COOH, $-(CH_2)_n$ -C(O)-COOH, tetrazole, -C(O)-NH₂, $-(CH_2)_n$ -C(O)-NH₂,






10 n is an integer from 0 to 3;

R_9 is selected from H, halogen, -CF₃, -OH, -COOH, $-(CH_2)_n$ -COOH, $-(CH_2)_n$ -C(O)-COOH, -C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -NH(C₁-C₆ alkyl), or -N(C₁-C₆ alkyl)₂;


15 n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, -CF₃, -OH, $-(CH_2)_n$ -COOH, $-(CH_2)_n$ -C(O)-COOH, -C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂,

20




- 14 -



n is an integer from 0 to 3;


15 R_{11} is selected from H, C_1-C_6 lower alkyl, C_1-C_6 cycloalkyl, $-CF_3$, $-COOH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$,

- 15 -



with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,


10



n is an integer from 0 to 3;

15 or a pharmaceutically acceptable salt thereof.

One preferred subset of the compounds of this invention, as defined above, are those having the formulae:



- 16 -

5

It will be understood in the group above that substituents R_3 and R_4 are bound to the indole or indoline ring's 2- or 3-position and the R_1 , R_1 , and R_2 groups are bound to one of the indole or indoline ring's 4-, 5-, 6- or 7-position carbon atoms.

10

One group of compounds within this invention are those in which the R_1 and R_3 groups are hydrogen and the substituents at the other indole or indoline positions are as described above.

15

Another group of this invention comprises compounds in which the R_1 and R_3 groups are hydrogen and the groups at R_1 , R_4 , and R_5 are as defined above. Within this group are two further preferred groups. In the first, R_1 is in the indole or indoline 5-position and in the second R_1 is in the indole or indoline 6 position.

20

In a further preferred group herein, R_1 is in the indole or indoline 5-position and is benzyloxy, R_2 is hydrogen and R_3 , R_4 and R_5 are as defined above.

25

In an another preferred group of this invention R_1 is in the indole or indoline 5 or 6 position and is cyclopentylcarboxamide or cyclopentyloxycarbonylamino, R_2 is hydrogen, and R_3 , R_4 and R_5 are as defined above.


30

A further preferred group of this invention consists of R_1 and R_2 at the indole or indoline 5 and or 6 position and are each selected from the group consisting of C_1 - C_6 alkoxy, cyano, sulfonyl and halo, R_2 is hydrogen, and R_3 , R_4 and R_5 are as defined above.

- 17 -

5 It is also understood that there is a further preferred subgroup within each of the groups set forth herein wherein the core molecule is an indole moiety, rather than an indoline. There is also understood to be a second group within each wherein the core molecule is an indoline moiety.

10 Preferred compounds of this invention include those of the following formulae:




wherein:

15 R₁ is selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NO₂, -NH₂, -HN(C₁-C₆), -N(C₁-C₆)₂, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, -CF₃, or -OH; or R₁ is a moiety of the formulae:

20

- 18 -



10 R_6 is selected from H, C_1-C_6 alkyl, C_1-C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NH_2$, $-NO_2$, $-CN$, $-CF_3$, or $-OH$;

15 R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1-C_6 alkyl, C_3-C_5 cycloalkyl, C_1-C_6 alkoxy, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, quinolyl, $(CH_2)_n$ phenyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, $-(CH_2)_n$ phenyl-O-phenyl, $-(CH_2)_n$ phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ phenyl-(O- CH_2 -phenyl) $_2$, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NH_2$, $-NO_2$, $-CF_3$, CO_2H , or $-OH$;

20


- 19 -

5 n is an integer from 0 to 3;

R₂ is selected from H, halogen, -CF₃, -OH, -C₁-C₁₀ alkyl, preferably -C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -CN, -NO₂, -NH₂, -NH-C₁-C₆ alkyl, -N(C₁-C₆ alkyl)₂,

10 -N-SO₂-C₁-C₆ alkyl, or -SO₂-C₁-C₆ alkyl;

R₃ is selected from H, -CF₃, -COOH, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, C₃-C₁₀ cycloalkyl, -C₁-C₆ alkyl-C₃-C₁₀ cycloalkyl, -CHO, halogen, or a moiety of the formulae:



wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₁-C₆ alkyl, C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl; pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents
 25 selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;


- 20 -

5

R_4 is selected from the group of C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, $-(CH_2)_n$ - C_3 - C_6 cycloalkyl, $-(CH_2)_n$ -S- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, $-(CH_2)_n$ -O- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, or the groups of:

10

a) $-(CH_2)_n$ -phenyl-O-phenyl, $-(CH_2)_n$ -phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ -phenyl-(O- CH_2 -phenyl)₂, or a moiety of the formulae:



15


wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C_3 - C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, -OH, $-C_1$ - C_6 alkyl, C_1 - C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O; or

20

b) a moiety of the formulae $-(CH_2)_n$ -A, $-(CH_2)_n$ -S-A, or $-(CH_2)_n$ -O-A, wherein A is the moiety:

25

- 21 -





5

wherein

D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;


10 B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, or -NO₂; or

c) a moiety of the formulae:



15

- 22 -



wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$; or

d) a moiety of the formula $-\text{L}^2\text{-M}^2$, wherein:

L^2 indicates a linking or bridging group of the formulae $-(\text{CH}_2)_n-$, $-\text{S}-$, $-\text{O}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-(\text{CH}_2)_n-$, $-(\text{CH}_2)_n-\text{O}-(\text{CH}_2)_n-$, or $-(\text{CH}_2)_n-\text{S}-$

$(\text{CH}_2)_n-$, $-\text{C}(\text{O})\text{C}(\text{O})\text{X}$;

where $\text{X} = \text{O}, \text{N}$

M^2 is selected from the group of $\text{C}_1\text{-C}_6$ lower alkyl, $\text{C}_1\text{-C}_6$ lower alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being

- 23 -

5 optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or


10 i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

15 ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH; or

25 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;

30 n is an integer from 0 to 3;



5 R₅ is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, -CH₂-phenyl-C(O)-benzothiazole, (CH₂)_n-CH=CH-COOH,





10

- 25 -






5



10



- 26 -




5

n is an integer from 0 to 3;

R_8 is selected from H, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, tetrazole, $-\text{C(O)}-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C(O)}-\text{NH}_2$,

10



n is an integer from 0 to 3;


R_9 is selected from H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$;

15 n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$,


- 27 -

- 5 $-(CH_2)_n-C(O)-COOH$, $-C_1-C_6$ alkyl, $-O-C_1-C_6$ alkyl, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂,




- 10 n is an integer from 0 to 3;

R₁₁ is selected from H, C₁-C₆ lower alkyl, -CF₃, -COOH, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, or




- 15 with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R₅, R₈, R₉, R₁₀, and/or R₁₁ shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,

- 28 -



n is an integer from 0 to 3;
 or a pharmaceutically acceptable salt thereof.

10 Another group of compounds of this invention have the following formulae:




wherein:

15 R_1 is selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6$ alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, phenyl, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or a moiety of the formulae:

20

- 29 -




- 10 R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;
- 15 R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N-(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH-(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-NH-(C_1-C_6 \text{ alkyl})$, $-N-(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, the pyridinyl, phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen,
- 20 C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;
n is an integer from 0 to 3;

10075079-021202

- 30 -

5 R_2 is selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_{10}$ alkyl, preferably $-C_1-C_6$ alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-CHO$, $-CN$, $-NO_2$, $-NH_2$, $-NH-C_1-C_6$ alkyl, $-N(C_1-C_6 \text{ alkyl})_2$, $-N-SO_2-C_1-C_6$ alkyl, or $-SO_2-C_1-C_6$ alkyl;

10 R_3 is selected from H, $-CF_3$, $-COOH$, C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, $-C_1-C_6$ alkyl- C_3-C_{10} cycloalkyl, $-CHO$, halogen, or a moiety of the formulae:




25

- 31 -

5 R_4 is selected from the group of C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, $-(CH_2)_n$ - C_3 - C_6 cycloalkyl, $-(CH_2)_n$ -S- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, $-(CH_2)_n$ -O- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, or the groups of:


10 a) $-(CH_2)_n$ -phenyl-O-phenyl, $-(CH_2)_n$ -phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ -phenyl-(O- CH_2 -phenyl)₂, $-CH_2$ -phenyl-C(O)-benzothiazole or a moiety of the formulae:



15 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C_3 - C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being
20 optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1$ - C_6 alkyl, C_1 - C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O; or

25 b) a moiety of the formulae $-(CH_2)_n$ -A, $-(CH_2)_n$ -S-A, or $-(CH_2)_n$ -O-A, wherein A is the moiety:

- 32 -




5

wherein

D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;


10 B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, or -NO₂; or

c) a moiety of the formulae:



15

- 33 -



5

wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, or -NO₂; or

10

d) a moiety of the formula -L²-M², wherein:

L² indicates a linking or bridging group of the formulae -(CH₂)_n-, -S-, -O-, -SO₂-, -C(O)-, -(CH₂)_n-C(O)-, -(CH₂)_n-C(O)-(CH₂)_n-, -(CH₂)_n-O-(CH₂)_n-, or -(CH₂)_n-S-(CH₂)_n-, -C(O)C(O)X;

15

where X = O, N

M² is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, C₃-C₁₀ cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being

- 34 -

5 optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

10 i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

15 ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH; or

20

25 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;


n is an integer from 0 to 3;

30




10075079-031203

- 35 -

- 5 R_5 is selected from $-\text{COOH}$, $-\text{C(O)}-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-\text{CH}_2\text{-phenyl-C(O)-benzothiazole}$,
 $(\text{CH}_2)_n-\text{CH=CH-COOH}$,




10








15

- 36 -

5




- 37 -



5

n is an integer from 0 to 3;

R_8 is selected from H, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, tetrazole, $-\text{C(O)}-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C(O)}-\text{NH}_2$,



10

n is an integer from 0 to 3;


R_9 is selected from H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$;

15 n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$,


- 38 -

5




10 n is an integer from 0 to 3;

R₁₁ is selected from H, C₁-C₆ lower alkyl, -CF₃, -COOH, -(CH₂)_n-COOH, -(CH₂)_n-C(O)-COOH, or



15 with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R₅, R₈, R₉, R₁₀, and/or R₁₁ shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: -C(O)-NH₂, -(CH₂)_n-C(O)-NH₂,

- 39 -



n is an integer from 0 to 3;

or a pharmaceutically acceptable salt thereof.

10 One group of compounds within this invention are those in which the indole or indoline 3-position (R₃) is substituted only by hydrogen and the substituents at the other indole or indoline positions are as described above.


15 Another group of this invention comprises compounds in which R₂ is hydrogen and the groups at R₁, R₃, R₄ and R₅ are as defined above. Within this group are two further preferred groups. In the first, R₁ is in the indole or indoline 5 position and in the second R₁ is in the indole or indoline 6 position.

20 In a further preferred group herein, R₁ is in the indole or indoline 5-position and is benzyloxy, optionally substituted as described above, R₂ is hydrogen and R₃, R₄ and R₅ are as defined above.

25 It is also understood that there is a further preferred subgroup within each of the groups set forth herein wherein the core molecule is an indole moiety, rather than an indoline. There is also understood to be a second group within each wherein the core molecule is an indoline moiety.

Another subset of compounds of this invention have the following formulae:


- 40 -



5

wherein:

R_1 is selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6$ alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, phenyl, $-O$ -phenyl, benzyl, $-O$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is a moiety of the formulae:



15

or

10075079-02100

- 41 -



5 R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;

10 R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N-(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH-(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-NH-(C_1-C_6 \text{ alkyl})$, $-N-(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, the pyridinyl, phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen,
15 C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;
 n is an integer from 0 to 3;


R_2 is selected from H, halogen, $-CF_3$, -OH, $-C_1$ - C_{10} alkyl, preferably $-C_1$ - C_6 alkyl, C_1 - C_{10} alkoxy, preferably C_1 - C_6 alkoxy, -CHO, -CN, $-NO_2$, $-NH_2$, $-NH-C_1-C_6$ alkyl, $-N(C_1-C_6 \text{ alkyl})_2$, $-N-SO_2-C_1-C_6$ alkyl, or $-SO_2-C_1-C_6$ alkyl;

20

R_3 is selected from H, $-CF_3$, $-COOH$, C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, C_3 - C_{10} cycloalkyl, $-C_1-C_6$ alkyl- C_3 - C_{10} cycloalkyl, -CHO, halogen, or a moiety of the formulae:






- 42 -



wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₁-C₆ alkyl, C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or
10 pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;

15 R₄ is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, -(CH₂)_n-C₃-C₆ cycloalkyl, -(CH₂)_n-S-(CH₂)_n-C₃-C₅ cycloalkyl, -(CH₂)_n-O-(CH₂)_n-C₃-C₅ cycloalkyl, or the groups of:


a) -(CH₂)_n-phenyl-O-phenyl, -(CH₂)_n-phenyl-CH₂-phenyl, -(CH₂)_n-O-phenyl-CH₂-phenyl,
20 -(CH₂)_n-phenyl-(O-CH₂-phenyl)₂, -CH₂-phenyl-C(O)-benzothiazole or a moiety of the formulae:



- 43 -

- 5 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring
10 containing one heteroatom selected from N, S, or O, preferably S or O; or

b) a moiety of the formulae -(CH₂)_n-A, -(CH₂)_n-S-A, or -(CH₂)_n-O-A, wherein A is the moiety:




- 15 wherein

D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;


B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy,
20 or -NO₂; or

c) a moiety of the formulae:



- 44 -

5



wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$; or

d) a moiety of the formula $-\text{L}^2\text{-M}^2$, wherein:

L^2 indicates a linking or bridging group of the formulae $-(\text{CH}_2)_n-$, $-\text{S}-$, $-\text{O}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-(\text{CH}_2)_n-$, $-(\text{CH}_2)_n-\text{O}-(\text{CH}_2)_n-$, or $-(\text{CH}_2)_n-\text{S}-(\text{CH}_2)_n-$, $-\text{C}(\text{O})\text{C}(\text{O})\text{X}$;
where X = O, N

- 45 -

5 M² is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, C₃-C₁₀ cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

10

i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected
15 from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring
20 being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH; or


25 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected
30 from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;

n is an integer from 0 to 3;



- 46 -

5



R_5 is selected from $-\text{COOH}$, $-\text{C(O)}-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-\text{CH}_2\text{-phenyl-C(O)-benzothiazole}$,
 $(\text{CH}_2)_n-\text{CH=CH-COOH}$,






10




15






- 48 -



5

n is an integer from 0 to 3;

R_8 is selected from H, ---COOH , $\text{---(CH}_2)_n\text{---COOH}$, $\text{---(CH}_2)_n\text{---C(O)---COOH}$, tetrazole, ---C(O)---NH_2 , $\text{---(CH}_2)_n\text{---C(O)---NH}_2$,



10

n is an integer from 0 to 3;



R_9 is selected from H, halogen, ---CF_3 , ---OH , $\text{---(CH}_2)_n\text{---COOH}$, $\text{---(CH}_2)_n\text{---C(O)---COOH}$, $\text{---C}_1\text{---C}_6$ alkyl, $\text{---O---C}_1\text{---C}_6$ alkyl, $\text{---NH(C}_1\text{---C}_6\text{ alkyl)}$, $\text{---N(C}_1\text{---C}_6\text{ alkyl)}_2$;

15 n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, ---CF_3 , ---OH , $\text{---(CH}_2)_n\text{---COOH}$,

- 49 -


5 $-(CH_2)_n-C(O)-COOH$, $-C_1-C_6$ alkyl, $-O-C_1-C_6$ alkyl, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂,



10

n is an integer from 0 to 3;


R_{11} is selected from H, C_1-C_6 lower alkyl, $-CF_3$, $-COOH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, or



15

with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,


- 50 -



n is an integer from 0 to 3;

or a pharmaceutically acceptable salt thereof.


Further preferred among the compounds of this invention are those having the
 10 formulae:



wherein:

15 R_1 is selected from halogen, $-NH_2$, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-N$ -benzyl, $-N$ -benzyl- O -phenyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is a moiety of the formulae:

- 51 -



R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-\text{NO}_2$, $-\text{CF}_3$, or $-\text{OH}$;


R_7 is selected from $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{N}-(\text{C}_1-\text{C}_6 \text{ alkyl})_2$, $-(\text{CH}_2)_n-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{CF}_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}-(\text{C}_1-\text{C}_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-

- 52 -


- 5 benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;

n is an integer from 0 to 3;

- 10 R_3 is selected from H, $-CF_3$, $-COOH$, C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, $-C_1-C_6$ alkyl- C_3-C_{10} cycloalkyl, $-CHO$, halogen, or a moiety of the formulae:



- 15 $\text{---} \text{O} \text{---} \text{(CH}_2\text{)}_n \text{---} \text{Y}$, $\text{---} \text{(CH}_2\text{)}_n \text{---} \text{O} \text{---} \text{Y}$,




- wherein n is independently selected in each appearance as an integer from 0 to 3,
 20 preferably 0 to 2, more preferably 0 to 1, Y is C_1-C_6 alkyl, C_3-C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6$ alkyl, C_1-C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O,
 25 preferably S or O;

- 53 -

5 R_4 is selected from the group of C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, $-(CH_2)_n$ - C_3 - C_6 cycloalkyl, $-(CH_2)_n$ -S- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, $-(CH_2)_n$ -O- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, or the groups of:


10 a) $-(CH_2)_n$ -phenyl-O-phenyl, $-(CH_2)_n$ -phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ -phenyl-(O- CH_2 -phenyl)₂, $-CH_2$ -phenyl-C(O)-benzothiazole or a moiety of the formulae:



20 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C_3 - C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1$ - C_6 alkyl, C_1 - C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O; or

25 b) a moiety of the formulae $-(CH_2)_n$ -A, $-(CH_2)_n$ -S-A, or $-(CH_2)_n$ -O-A, wherein A is the moiety:

- 54 -



5


wherein

D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;

B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, or -NO₂; or

10


c) a moiety of the formulae:



15

10075079.034700

- 55 -



wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$; or

d) a moiety of the formula $-\text{L}^2\text{-M}^2$, wherein:

L^2 indicates a linking or bridging group of the formulae $-(\text{CH}_2)_n-$, $-\text{S}-$, $-\text{O}-$, $-\text{SO}_2-$, $-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-$, $-(\text{CH}_2)_n-\text{C}(\text{O})-(\text{CH}_2)_n-$, $-(\text{CH}_2)_n-\text{O}-(\text{CH}_2)_n-$, or $-(\text{CH}_2)_n-\text{S}-(\text{CH}_2)_n-$, $-\text{C}(\text{O})\text{C}(\text{O})\text{X}$;
 where X = O, N

M^2 is selected from the group of $\text{C}_1\text{-C}_6$ lower alkyl, $\text{C}_1\text{-C}_6$ lower alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being

- 56 -

5 optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

10 i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or





15 ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH; or

25 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;



30 R_s is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, -CH₂-phenyl-C(O)-benzothiazole,

- 57 -




5 $(\text{CH}_2)_n\text{-CH=CH-COOH}$,



10




- 58 -



10376379.042201


- 59 -



5

n is an integer from 0 to 3;

R₈ is selected from H, -COOH, -(CH₂)_n-COOH, -(CH₂)_n-C(O)-COOH, tetrazole, -C(O)-NH₂, -(CH₂)_n-C(O)-NH₂,



n is an integer from 0 to 3;






R₉ is selected from H, halogen, -CF₃, -OH, -(CH₂)_n-COOH, -(CH₂)_n-C(O)-COOH, -C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂;

15 n is an integer from 0 to 3;

R₁₀ is selected from the group of H, halogen, -CF₃, -OH, -(CH₂)_n-COOH,

- 60 -


- 5 $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$,



10

n is an integer from 0 to 3;


R_{11} is selected from H, C_1-C_6 lower alkyl, $-\text{CF}_3$, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, or



15

with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-\text{C}(\text{O})-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NH}_2$,


- 61 -



n is an integer from 0 to 3;

or a pharmaceutically acceptable salt thereof.

Further preferred among the compounds of this invention are those having the
 10 formulae:




wherein:

R_1 is selected from halogen, $-NH_2$, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-N$ -benzyl, $-N$ -benzyl- O -phenyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is or a moiety of the formulae:

15

- 62 -



R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by
 15 from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;


R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N-(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH-(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-NH-(C_1-C_6 \text{ alkyl})$, $-N-(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-
 20

- 63 -

- 5 benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;

n is an integer from 0 to 3;

- 10 R_3 is selected from H, halogen, $-CF_3$, $-OH$, C_1-C_6 alkyl, C_1-C_6 alkoxy, benzyl, benzyloxy, phenyl, phenyloxy, $-C(O)$ -phenyl, $-C(O)$ -benzyl, $-CH_2-(C_3-C_6 \text{ cycloalkyl})$, $-C(O)-OH$, $-CH=O$, $-C(O)-C_1-C_6$ alkyl, $-C(O)-O-C_1-C_6$ alkyl, $-C(O)-CF_3$, $-(CH_2)_n-S-CH_2-(C_3-C_6 \text{ cycloalkyl})$,



15

the rings of the relevant R_3 groups being optionally substituted by from 1 to 3 groups selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-CF_3$, $-C(O)-OH$, or $-OH$;


n is an integer from 0 to 3;

20

R_4 is selected from the group of C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, $-(CH_2)_n-C_3-C_6 \text{ cycloalkyl}$, $-(CH_2)_n-S-(CH_2)_n-C_3-C_5 \text{ cycloalkyl}$, $-(CH_2)_n-O-(CH_2)_n-C_3-C_5 \text{ cycloalkyl}$, or the groups of:


- 25 a) $-(CH_2)_n$ -phenyl-O-phenyl, $-(CH_2)_n$ -phenyl- CH_2 -phenyl, $-(CH_2)_n$ -O-phenyl- CH_2 -phenyl, $-(CH_2)_n$ -phenyl-(O- CH_2 -phenyl) $_2$, $-CH_2$ -phenyl- $C(O)$ -benzothiazole or a moiety of the formulae:

- 64 -



- 10 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring
- 15 containing one heteroatom selected from N, S, or O, preferably S or O; or

b) a moiety of the formulae -(CH₂)_n-A, -(CH₂)_n-S-A, or -(CH₂)_n-O-A, wherein A is the moiety:



- 20 wherein


D is H, C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, or -CF₃;

- B and C are independently selected from phenyl, pyridinyl, furyl, thienyl, pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably 1 to 2, substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy,
- 25 or -NO₂; or

- 65 -

5

c) a moiety of the formulae:



10

wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$; or

- 66 -

5 d) a moiety of the formula $-L^2-M^2$, wherein:

L^2 indicates a linking or bridging group of the formulae $-(CH_2)_n-$, $-S-$, $-O-$, $-SO_2-$, $-C(O)-$, $-(CH_2)_n-C(O)-$, $-(CH_2)_n-C(O)-(CH_2)_n-$, $-(CH_2)_n-O-(CH_2)_n-$, or $-(CH_2)_n-S-(CH_2)_n-$, $-C(O)C(O)X$;

10 where $X = O, N$

M^2 is selected from the group of C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, or $-CF_3$; or

20 i) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, or $-CF_3$; or

25 ii) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to pyridine, pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_{10} alkyl, preferably C_1-C_6 alkyl, C_1-C_{10} alkoxy, preferably C_1-C_6 alkoxy, $-CHO$, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$ or $-OH$; or


30

- 67 -

- 5 iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_{10} alkyl, preferably C_1 - C_6 alkyl, C_1 - C_{10} alkoxy, preferably C_1 - C_6 alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;
 10 n is an integer from 0 to 3;


R_5 is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, -CH₂-phenyl-C(O)-benzothiazole,

- 15 (CH₂)_n-CH=CH-COOH,






- 68 -






5



10




- 69 -



5

n is an integer from 0 to 3;

R_8 is selected from H, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, tetrazole, $-\text{C}(\text{O})-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NH}_2$,



10

n is an integer from 0 to 3;

R_9 is selected from H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$;




15

n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$,

- 70 -


- 5 $-(CH_2)_n-C(O)-COOH$, $-C_1-C_6$ alkyl, $-O-C_1-C_6$ alkyl, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂,



10

n is an integer from 0 to 3;


R_{11} is selected from H, C_1-C_6 lower alkyl, $-CF_3$, $-COOH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, or



15

with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,


- 71 -



5





n is an integer from 0 to 3;

- 10 Further preferred among the compounds of this invention are those having the formulae:




wherein:

- 15 R_1 is selected from Halogen, $-NH_2$, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-N$ -benzyl, $-N$ -benzyl- O -phenyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is or a moiety of the formulae:



20

- 72 -




R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by
10 from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;

R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N-(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH-(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1-C_6 alkyl, C_3-C_5 cycloalkyl, C_1-C_6 alkoxy, $-NH-(C_1-C_6 \text{ alkyl})$, $-N-(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally
15 substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;

n is an integer from 0 to 3;


20

R_3 is selected from H, $-CF_3$, $-COOH$, C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, $-C_1-C_6 \text{ alkyl}-C_3-C_{10} \text{ cycloalkyl}$, $-CHO$, halogen, or a moiety of the
formulae:



- 73 -

5




- 10 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₁-C₆ alkyl, C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five
15 membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;


R₄ is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, -(CH₂)_n-C₃-C₆ cycloalkyl, -(CH₂)_n-S-(CH₂)_n-C₃-C₅ cycloalkyl, -(CH₂)_n-O-(CH₂)_n-C₃-C₅ cycloalkyl, or the groups of:

20

- a) -(CH₂)_n-phenyl-O-phenyl, -(CH₂)_n-phenyl-CH₂-phenyl, -(CH₂)_n-O-phenyl-CH₂-phenyl, -(CH₂)_n-phenyl-(O-CH₂-phenyl)₂, -CH₂-phenyl-C(O)-benzothiazole or a moiety of the formulae:




25



- 74 -



5



wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C₃-C₅ cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl or pyrrolyl; rings of these groups being
10 optionally substituted by from 1 to 3 substituents selected from H, halogen, -CF₃, -OH, -C₁-C₆ alkyl, C₁-C₆ alkoxy, -NH₂, -NO₂ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;







n is an integer from 0 to 3;

R₅ is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, -
15 CH₂-phenyl-C(O)-benzothiazole,
(CH₂)_n-CH=CH-COOH,



20

- 75 -



10

n is an integer from 0 to 3;

R_8 is selected from H, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, tetrazole, $-\text{C}(\text{O})-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NH}_2$,

n is an integer from 0 to 3;


15

- 76 -

- 5 R_9 is selected from H, halogen, $-CF_3$, $-OH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, $-C_1-C_6$ alkyl, $-O-C_1-C_6$ alkyl, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂;

n is an integer from 0 to 3;


- 10 R_{10} is selected from the group of H, halogen, $-CF_3$, $-OH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, $-C_1-C_6$ alkyl, $-O-C_1-C_6$ alkyl, $-NH(C_1-C_6$ alkyl), $-N(C_1-C_6$ alkyl)₂,



15


n is an integer from 0 to 3;

R_{11} is selected from H, C_1-C_6 lower alkyl, $-CF_3$, $-COOH$, $-(CH_2)_n-COOH$, $-(CH_2)_n-C(O)-COOH$, or




- 77 -

- 5 with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,




- 10 n is an integer from 0 to 3;
 Further preferred among the compounds of this invention are those having the formulae:



- 15 wherein:
 R_1 is selected from halogen, $-NH_2$, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-N$ -benzyl, $-N$ -benzyl- O -phenyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is or a moiety of the formulae:

20

- 78 -



R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-\text{NO}_2$, $-\text{CF}_3$, or $-\text{OH}$;

15


R_7 is selected from $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{N}-(\text{C}_1-\text{C}_6 \text{ alkyl})_2$, $-(\text{CH}_2)_n-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{CF}_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-\text{NH}-(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}-(\text{C}_1-\text{C}_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally

- 79 -

5 substituted by from 1 to 3 substituents selected from halogen, C_1-C_6 alkyl, C_1-C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;

n is an integer from 0 to 3;

R_3 is selected from H, $-CF_3$, $-COOH$, C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, C_3-C_{10} cycloalkyl, $-C_1-C_6$ alkyl- C_3-C_{10} cycloalkyl, $-CHO$, halogen, or a moiety of the
 10 formulae:




15

wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C_1-C_6 alkyl, C_3-C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or
 20 pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6$ alkyl, C_1-C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;

R_4 is selected from the group of C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, $-(CH_2)_n-$
 25 C_3-C_6 cycloalkyl, $-(CH_2)_n-S-(CH_2)_n-C_3-C_5$ cycloalkyl, $-(CH_2)_n-O-(CH_2)_n-C_3-C_5$ cycloalkyl, or the groups of:

- 80 -

- 5 a) a moiety of the formulae $-(CH_2)_n-A$, $-(CH_2)_n-S-A$, or $-(CH_2)_n-O-A$,
 wherein A is the moiety:




wherein

D is H, C_1-C_6 lower alkyl, C_1-C_6 lower alkoxy, or $-CF_3$;

- 10 B and C are independently selected from phenyl, pyridinyl, furyl, thienyl,
 pyrimidinyl or pyrrolyl groups, each optionally substituted by from 1 to 3, preferably
 1 to 2, substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6$ alkyl, C_1-C_6 alkoxy,
 or $-NO_2$;


R_5 is selected from $-COOH$, $-C(O)-COOH$, $-(CH_2)_n-C(O)-COOH$, $-(CH_2)_n-COOH$, -

- 15 CH_2 -phenyl- $C(O)$ -benzothiazole,
 $(CH_2)_n-CH=CH-COOH$,




- 81 -

5



10



n is an integer from 0 to 3;

R₈ is selected from H, -COOH, -(CH₂)_n-COOH, -(CH₂)_n-C(O)-COOH, tetrazole, -
 15 C(O)-NH₂, -(CH₂)_n-C(O)-NH₂,


- 82 -

5 n is an integer from 0 to 3;

R_9 is selected from H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$;

10 n is an integer from 0 to 3;


R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$,



n is an integer from 0 to 3;

20 R_{11} is selected from H, C_1-C_6 lower alkyl, $-\text{CF}_3$, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, or


- 83 -



5

with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-C(O)-NH_2$, $-(CH_2)_n-C(O)-NH_2$,

10




n is an integer from 0 to 3;

or a pharmaceutically acceptable salt thereof.

15

Further preferred among the compounds of this invention are those having the formulae:





20 wherein:



R_1 is selected from halogen, $-NH_2$, $-O$ -phenyl, benzyl, $-O$ -benzyl, $-N$ -benzyl, $-N$ -benzyl- O -phenyl, $-S$ -benzyl, the phenyl and benzyl rings of these groups being

- 84 -

- 5 optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-NH_2$, $-CN$, $-CF_3$, or $-OH$; or R_1 is or a moiety of the formulae:



10



15

R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;


20

- 85 -

- 5 R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N-(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH-(C_1-C_6 \text{ alkyl})$, $-CF_3$, $C_1-C_6 \text{ alkyl}$, $C_3-C_5 \text{ cycloalkyl}$, $C_1-C_6 \text{ alkoxy}$, $-NH-(C_1-C_6 \text{ alkyl})$, $-N-(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, $-O$ -phenyl, benzyl, $-O$ -benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, $C_1-C_6 \text{ alkyl}$, C_1-C_6
 10 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;

n is an integer from 0 to 3;

- 15 R_3 is selected from H, $-CF_3$, $-COOH$, $C_1-C_6 \text{ lower alkyl}$, $C_1-C_6 \text{ lower alkoxy}$, $C_3-C_{10} \text{ cycloalkyl}$, $-C_1-C_6 \text{ alkyl}-C_3-C_{10} \text{ cycloalkyl}$, $-CHO$, halogen, or a moiety of the formulae:









- 20 wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is $C_1-C_6 \text{ alkyl}$, $C_3-C_5 \text{ cycloalkyl}$, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents
 25 selected from H, halogen, $-CF_3$, $-OH$, $-C_1-C_6 \text{ alkyl}$, $C_1-C_6 \text{ alkoxy}$, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;

- 86 -

- 5 R_4 is selected from the group of C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, $-(CH_2)_n$ - C_3 - C_6 cycloalkyl, $-(CH_2)_n$ -S- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, $-(CH_2)_n$ -O- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, or the groups of:

a) a moiety of the formulae:

10



or

- 87 -



5 wherein Z is O or S and the phenyl and pyrimidinyl rings of each moiety are optionally and independently substituted by from 1 to 3 substituents selected from halogen, $-\text{CF}_3$, $-\text{OH}$, $-\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_6$ alkoxy, $-\text{NH}_2$, or $-\text{NO}_2$;

n is an integer from 0 to 3;


R_5 is selected from $-\text{COOH}$, $-\text{C}(\text{O})-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, -

10 $\text{CH}_2\text{-phenyl-C}(\text{O})\text{-benzothiazole}$,


$(\text{CH}_2)_n\text{-CH=CH-COOH}$,



15



- 88 -



5

n is an integer from 0 to 3;

R_8 is selected from H, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, tetrazole, $-\text{C(O)}-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C(O)}-\text{NH}_2$,

n is an integer from 0 to 3;

10


R_9 is selected from H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6$ alkyl), $-\text{N}(\text{C}_1-\text{C}_6$ alkyl)₂;

n is an integer from 0 to 3;

15


R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C(O)}-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6$ alkyl), $-\text{N}(\text{C}_1-\text{C}_6$ alkyl)₂,

- 89 -



n is an integer from 0 to 3;


10 R_{11} is selected from H, $\text{C}_1\text{-C}_6$ lower alkyl, $-\text{CF}_3$, $-\text{COOH}$, $-(\text{CH}_2)_n\text{-COOH}$, $-(\text{CH}_2)_n\text{-C(O)-COOH}$, or



15 with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-\text{C(O)-NH}_2$, $-(\text{CH}_2)_n\text{-C(O)-NH}_2$,


- 90 -

5



n is an integer from 0 to 3;




- 10 Further preferred among the compounds of this invention are those having the formulae:



wherein:

- 15 R₁ is selected from halogen, -NH₂, -O-phenyl, benzyl, -O-benzyl, -N-benzyl, -N-benzyl-O-phenyl, -S-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, -CF₃, or -OH; or R₁ is or a moiety of the formulae:

- 91 -



R_6 is selected from H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, -O-phenyl, benzyl, -O-benzyl, the phenyl and benzyl rings of these groups being optionally substituted by
 15 from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or -OH;


R_7 is selected from $-(CH_2)_n-COOH$, $-(CH_2)_n-N(C_1-C_6 \text{ alkyl})_2$, $-(CH_2)_n-NH(C_1-C_6 \text{ alkyl})$, $-CF_3$, C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, C_1 - C_6 alkoxy, $-NH(C_1-C_6 \text{ alkyl})$, $-N(C_1-C_6 \text{ alkyl})_2$, pyridinyl, thienyl, furyl, pyrrolyl, phenyl, -O-phenyl, benzyl, -O-
 20

- 92 -

- 5 benzyl, adamantyl, or morpholinyl, the rings of these groups being optionally substituted by from 1 to 3 substituents selected from halogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, $-NO_2$, $-CF_3$, or $-OH$;

n is an integer from 0 to 3;

- 10 R_3 is selected from H, $-CF_3$, $-COOH$, C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, C_3 - C_{10} cycloalkyl, $-C_1$ - C_6 alkyl- C_3 - C_{10} cycloalkyl, $-CHO$, halogen, or a moiety of the formulae:



- wherein n is independently selected in each appearance as an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1, Y is C_1 - C_6 alkyl, C_3 - C_5 cycloalkyl, phenyl, benzyl, naphthyl, pyridinyl, quinolyl, furyl, thienyl, morpholinyl, pyrrolidinyl, or pyrrolyl; rings of these groups being optionally substituted by from 1 to 3 substituents selected from H, halogen, $-CF_3$, $-OH$, $-C_1$ - C_6 alkyl, C_1 - C_6 alkoxy, $-NH_2$, $-NO_2$ or a five membered heterocyclic ring containing one heteroatom selected from N, S, or O, preferably S or O;

R_4 is selected from the group of C_1 - C_6 lower alkyl, C_1 - C_6 lower alkoxy, $-(CH_2)_n$ - C_3 - C_6 cycloalkyl, $-(CH_2)_n$ -S- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, $-(CH_2)_n$ -O- $(CH_2)_n$ - C_3 - C_5 cycloalkyl, or the groups of:

- 93 -

5

d) a moiety of the formula $-L^2-M^2$, wherein:

L^2 indicates a linking or bridging group of the formulae $-(CH_2)_n-$, $-S-$, $-O-$,
-SO₂-, $-C(O)-$, $-(CH_2)_n-C(O)-$, $-(CH_2)_n-C(O)-(CH_2)_n-$, $-(CH_2)_n-O-(CH_2)_n-$, or $-(CH_2)_n-S-$
10 $(CH_2)_n-C(O)C(O)X$;
where X = O,N

M^2 is selected from the group of C₁-C₆ lower alkyl, C₁-C₆ lower alkoxy, C₃-
C₁₀ cycloalkyl, phenyl or benzyl, the cycloalkyl, phenyl or benzyl rings being
15 optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl,
preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -NO₂, -NH₂, -CN, or -
CF₃; or

i) a five-membered heterocyclic ring containing one or two ring
20 heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole,
thiophene, imidazole, pyrazole, pyrrolidine, or tetrazole, the five-membered
heterocyclic ring being optionally substituted by from 1 to 3 substituents selected
from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆
alkoxy, -NO₂, -NH₂, -CN, or -CF₃; or

25


ii) a six-membered heterocyclic ring containing one, two or three ring
heteroatoms selected from N, S or O including, but not limited to pyridine,
pyrimidine, piperidine, piperazine, or morpholine, the six-membered heterocyclic ring
being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀
30 alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -
NH₂, -CN, -CF₃ or -OH; or

- 94 -

5








- iii) a bicyclic ring moiety containing from 8 to 10 ring atoms and optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, indole, indoline, naphthalene, purine, or quinoline, the bicyclic ring moiety being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -NO₂, -NH₂, -CN, -CF₃ or -OH;
- 10 n is an integer from 0 to 3;

- R₅ is selected from -COOH, -C(O)-COOH, -(CH₂)_n-C(O)-COOH, -(CH₂)_n-COOH, -CH₂-phenyl-C(O)-benzothiazole,
- 15 (CH₂)_n-CH=CH-COOH,



20

- 95 -



10 n is an integer from 0 to 3;

R₈ is selected from H, -COOH, -(CH₂)_n-COOH, -(CH₂)_n-C(O)-COOH, tetrazole, -C(O)-NH₂, -(CH₂)_n-C(O)-NH₂,

n is an integer from 0 to 3;


R₉ is selected from H, halogen, -CF₃, -OH, -(CH₂)_n-COOH,

15 -(CH₂)_n-C(O)-COOH, -C₁-C₆ alkyl, -O-C₁-C₆ alkyl, -NH(C₁-C₆ alkyl), -N(C₁-C₆ alkyl)₂;

- 96 -


5 n is an integer from 0 to 3;

R_{10} is selected from the group of H, halogen, $-\text{CF}_3$, $-\text{OH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, $-\text{C}_1-\text{C}_6$ alkyl, $-\text{O}-\text{C}_1-\text{C}_6$ alkyl, $-\text{NH}(\text{C}_1-\text{C}_6 \text{ alkyl})$, $-\text{N}(\text{C}_1-\text{C}_6 \text{ alkyl})_2$,




n is an integer from 0 to 3;

15 R_{11} is selected from H, C_1-C_6 lower alkyl, $-\text{CF}_3$, $-\text{COOH}$, $-(\text{CH}_2)_n-\text{COOH}$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{COOH}$, or



20 with a proviso that the complete moiety at the indole or indoline 1-position created by any combination of R_5 , R_8 , R_9 , R_{10} , and/or R_{11} shall contain at least one acidic moiety selected from or containing a carboxylic acid, a tetrazole, or a moiety of the formulae: $-\text{C}(\text{O})-\text{NH}_2$, $-(\text{CH}_2)_n-\text{C}(\text{O})-\text{NH}_2$,

- 97 -



5

n is an integer from 0 to 3;
or a pharmaceutically acceptable salt thereof.

Detailed Description of Preferred Embodiments

10

As used herein, the terms "aryl" and "substituted aryl" are understood to include monocyclic, particularly including five- and six-membered monocyclic, aromatic and heteroaromatic ring moieties and bicyclic aromatic and heteroaromatic ring moieties, particularly including those having from 9 to 10 ring atoms. Among these aryl groups are understood to be phenyl rings, including those found in phenoxy, benzyl, benzyloxy, biphenyl and other such moieties. The aryl and heteroaryl groups of this invention also include the following:

15

a) a five-membered heterocyclic ring containing one or two ring heteroatoms selected from N, S or O including, but not limited to, furan, pyrrole, thiophene, imidazole, pyrazole, isothiazole, isoxazole, pyrrolidine, pyrroline, imidazolidine, pyrazolidine, pyrazole, pyrazoline, imidazole, tetrazole, or oxathiazole; or

20

b) a six-membered heterocyclic ring containing one, two or three ring heteroatoms selected from N, S or O including, but not limited to, pyran, pyridine, pyrazine, pyrimidine, pyridazine, piperidine, piperazine, tetrazine, thiazine, thiadizine, oxazine, or morpholine; or

25

5

c) a bicyclic ring moiety optionally containing from 1 to 3 ring heteroatoms selected from N, S or O including, but not limited to benzofuran, chromene, indole, isoindole, indoline, isoindoline, naphthalene, purine, indolizine, indazole, quinoline, isoquinoline, quinolizine, quinazoline, cinnoline, phthalazine, or naphthyridine.

10

The "substituted aryl" groups of this invention include such moieties being optionally substituted by from 1 to 3 substituents selected from halogen, C₁-C₁₀ alkyl, preferably C₁-C₆ alkyl, C₁-C₁₀ alkoxy, preferably C₁-C₆ alkoxy, -CHO, -COOH or esters thereof, -NO₂, -NH₂, -CN, -CF₃ or -OH or combinations thereof, such as -CH₂CF₃, -NH(CH₃), etc.

15

A preferred subset of these groups, optionally substituted as just described, include moieties formed from benzene, pyridine, naphthylene or quinoline rings. A further preferred group includes those of furan, pyrrole, thiophene, pyrimidine, and morpholine rings. A preferred group of bicyclic aromatic groups includes benzofuran, indole, naphthalene, and quinoline rings.

20

The alkyl, alkenyl and alkynyl groups referred to herein indicate such groups having from 1 to 10, preferably 1 to 6 carbon atoms, and may be straight, branched or cyclic. Unless indicated otherwise, it is preferred that these groups be straight or branched. Halogens herein are understood to include F, Cl, Br and I.

25

The integers depicted by the variable n are in each appearance independently an integer from 0 to 3, preferably 0 to 2, more preferably 0 to 1. It will be understood the appearance of n in any instance is not definitive or limiting of the integer which

30

- 99 -

5 may be indicated by n elsewhere within the definition of the present invention. For instance, each of the integers indicated by n in the moiety $-(CH_2)_n-O-(CH_2)_n-$ may be selected from 0, 1, 2 or 3.

10 As used herein, "phospholipase enzyme activity" means positive activity in an assay for metabolism of phospholipids (preferably one of the assays described in Example 86 below). A compound has "phospholipase enzyme inhibiting activity" when it inhibits the activity of a phospholipase (preferably cPLA₂) in any available assay (preferably an assay described below in Example 86 or Example 87) for enzyme activity. In preferred embodiments, a compound has (1) an IC₅₀ value of less
15 than about 25 μM, preferably less than about 6 μM, in the LysoPC assay; (2) an IC₅₀ value of less than about 50 μM in the vesicle assay; (3) an IC₅₀ value of less than about 1 μM in the PMN assay; (4) an IC₅₀ value of less than about 15 μM in the Coumarine assay; and/or (5) measurable activity (preferably at least about 5% reduction in edema, more preferably at least about 10% reduction, more preferably at
20 least about 15%, most preferably 20-30%) in the rat carrageenan-induced footpad edema test.

Compounds of the present invention are useful for inhibiting phospholipase enzyme (preferably cPLA₂) activity and, therefore, are useful in "treating" (i.e.,
25 treating, preventing or ameliorating) inflammatory or inflammation-related responses or conditions (e.g., rheumatoid arthritis, psoriasis, asthma, inflammatory bowel disease, and other diseases mediated by prostaglandins, leukotrienes or PAF) and other conditions, such as osteoporosis, colitis, myelogenous leukemia, diabetes, wasting and atherosclerosis.

30

- 100 -

5 The present invention encompasses both pharmaceutical compositions and therapeutic methods of treatment or use which employ compounds of the present invention.

10 Compounds of the present invention may be used in a pharmaceutical composition when combined with a pharmaceutically acceptable carrier. Such a composition may also contain (in addition to a compound or compounds of the present invention and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition may further contain other anti-inflammatory agents. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with compounds of the present invention, or to minimize side effects caused by the compound of the present invention.

20

25 The pharmaceutical composition of the invention may be in the form of a liposome in which compounds of the present invention are combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323, all of which are incorporated herein by reference.

30

As used herein, the terms “pharmaceutically effective amount” or “therapeutically effective amount” means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, i.e., treatment, healing, prevention or amelioration of an inflammatory response or condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination, the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of a compound of the present invention is administered to a mammal having a condition to be treated. Compounds of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing other anti-inflammatory agents, cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more other anti-inflammatory agents, cytokines, lymphokines or other hematopoietic factors, compounds of the present invention may be administered either simultaneously with the other anti-inflammatory agent(s), cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering compounds of the present invention in combination with other anti-inflammatory agent(s), cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

- 102 -

5

Administration of compounds of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection.

10

When a therapeutically effective amount of compounds of the present invention is administered orally, compounds of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% compound of the present invention, and preferably from about 25 to 90% compound of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of compound of the present invention, and preferably from about 1 to 50% compound of the present invention.

25

When a therapeutically effective amount of compounds of the present invention is administered by intravenous, cutaneous or subcutaneous injection, compounds of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein solutions, having due regard to pH, isotonicity, stability, and the like, is within the

30

- 103 -

5 skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous,
or subcutaneous injection should contain, in addition to compounds of the present
invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection,
Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's
Injection, or other vehicle as known in the art. The pharmaceutical composition of
-10 the present invention may also contain stabilizers, preservatives, buffers, antioxidants,
or other additives known to those of skill in the art.

202120-62052001
15 The amount of compound(s) of the present invention in the pharmaceutical
composition of the present invention will depend upon the nature and severity of the
condition being treated, and on the nature of prior treatments which the patient has
undergone. Ultimately, the attending physician will decide the amount of compound
of the present invention with which to treat each individual patient. Initially, the
attending physician will administer low doses of compound of the present invention
and observe the patient's response. Larger doses of compounds of the present
20 invention may be administered until the optimal therapeutic effect is obtained for the
patient, and at that point the dosage is not increased further. It is contemplated that
the various pharmaceutical compositions used to practice the method of the present
invention should contain about 0.1 μ g to about 100 mg (preferably about .1 mg to
about 50 mg, more preferably about 1 mg to about 2 mg) of compound of the present
25 invention per kg body weight.

The duration of intravenous therapy using the pharmaceutical composition of
the present invention will vary, depending on the severity of the disease being treated
and the condition and potential idiosyncratic response of each individual patient. It is
30 contemplated that the duration of each application of the compounds of the present
invention will be in the range of 12 to 24 hours of continuous intravenous

- 104 -

5 administration. Ultimately the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

Compounds of the present invention can be made according to the methods
-10 and examples described below. Synthesis of preferred compounds of the present invention are described in the examples below.

Method A

Ethyl 5-nitro-2-carboxylate indole is chlorinated in the 3-position by the agency of N-
15 chlorosuccinamide in a solvent such as DMF or DMSO at an elevated temperature of 40°C - 80°C. The ester is then reduced in a three step procedure. First the ester is hydrolysed under basic conditions with a base such as sodium hydroxide or potassium hydroxide in a solvent system such as water:methanol:THF. The acid is then activated by the addition of carbonyl diimidazole in THF and reduced with a reducing
20 agent such as sodium borohydride or sodium triacetoxyborohydride in an alcoholic solvent system such as methanol or ethanol. The resulting alcohol is protected as the TBDMS ether with TBDMSCl in a solvent such as DMF, methylene chloride or THF with a base such as triethylamine or imidazole. The indole nitrogen is then alkylated with methyl 4-bromomethylbenzoate in a solvent such as THF, acetonitrile or DMF
25 with a base such as sodium hydride, *n*-BuLi or potassium bis(trimethylsilyl)amide. The 5-nitro group is then reduced by exposure to H₂ in the presence of a catalyst such as Pt/C or Pd/C in a solvent such as ethyl acetate, methanol or THF or a mixture of two or all of the three. The amine is then acylated with cyclopentylcarbonyl chloride in a biphasic system of saturated sodium bicarbonate and methylene chloride. R₁ is
30 then introduced in a two step procedure wherein the TBDMS ether is converted to a bromide by exposure to dibromotriphenylphosphorane in methylene chloride and then

- 105 -

5 the crude bromide is displaced by a variety of thiols or phenols in a solvent such as THF, methylene chloride or DMF in the presence of a base such as potassium carbonate or cesium carbonate. The product esters are then prepared by the hydrolysis of the ester under basic conditions with sodium hydroxide in a solvent system such as water:MeOH:THF.

10

Method B

15 Acylation at the 3-position of indole I with an acylating agent such as naphthoyl chloride could be accomplished using ethylmagnesium bromide in a solvent such as THF to give II. Alkylation of the indole nitrogen could be accomplished by exposure to a suitable base such as sodium hydride followed by treatment with the appropriate. Deprotection of the hydroxy protecting group with tetrabutylammonium fluoride and oxidation with a suitable oxidizing agent provided IV. A Horner-Wittig reaction with trimethoxyphosphonoacetate in a suitable solvent such as tetrahydrofuran gave the unsaturated ester V which could be deprotected at the indole 1-position with a
20 suitable reagent system such as hydrofluoric acid in acetonitrile. Saponification of the remaining acid group gave the compound VI

Method C

25 Indole I can be converted to II in two steps: (1) reduction with LAH in a solvent such as THF and (2) silylation with t-butyldimethylsilyl chloride (TBDMSCl) in a solvent such as dichloromethane or DMF in the presence of a base such as imidazole. Treatment of II with Grignard reagent such as ethyl magnesium bromide in a solvent such as THF at -60°C, acylation of the resulting magnesium salt with a suitable acyl chloride such as acetyl chloride in ether and finally, alkylation on the nitrogen with an
30 alkyl halide such as methyl(4-bromomethyl)benzoate in the presence of a strong base such as NaH in DMF affords ketone III. The silyl group on III is removed using

- 106 -

5 tetrabutylammonium fluoride in a solvent such as THF, the resulting alcohol is then
converted to bromide using carbon tetrabromide and bis(diphenylphosphino)ethane in
a solvent such as dichloromethane to yield bromide IV. Displacement of the bromine
of IV with a thiol compound in the presence of a base such as cesium carbonate, or
with an alcohol in the presence of a strong base such as NaH in DMF affords V
-10 (sulfide or ether respectively).

Method D

The protected alcohol was deprotected in a suitable solvent such as THF and the
resulting alcohol was functionalized to a halide using carbon tetrabromide or
15 methanesulfonyl chloride and then reacted with an oxygen nucleophile, coupled with
prior deprotonation with a strong base such as sodium hydride, or a sulfur
nucleophile in the presence of cesium carbonate in DMF or THF. The nitro group
could then be reduced to an amine via a Pt/carbon hydrogenation protocol or copper
acetate sodium borohydride procedure. The resulting amine could be
20 hydrolyzed, using a standard procedure of sodium hydroxide in THF/MeOH or
coupled to a variety of acylating reagents, such as acid chlorides, chloroformates and
isocyanates where the reactions are generally performed in the presence of a base in a
solvent such as THF or dichloromethane. The amine could also be acylated via an
EDCI coupling procedure with a variety of acids. The starting amine could also be
25 alkylated by a reductive amination procedure using a variety of aldehydes and sodium
triacetoxymethylborohydride as the reducing agent. These functionalized amines could be
hydrolyzed to yield the desired acids which could also be converted to the
acylsulfonamide by EDCI coupling with a sulfonamide. Alternatively, the
functionalized amines could be alkylated further by reaction with a strong base and an
30 alkyl halide and then hydrolyzed under the standard conditions to yield the requisite
product.

- 107 -


5

Method E

The starting indole, with or without C2 substitution, was functionalized at C3 by using DMF/POCl₃ conditions or the magnesium salt of the indole was acylated with a variety of acid chlorides to form the ketones. These products were then N-alkylated by the action of a strong base and a variety of alkyl or aryl halo esters. When R' is a nitro group, at this time the nitro was reduced with Pt/C and H₂ or copper acetate and sodium borohydride to the amine which was then acylated with a variety of acid chlorides, isocyanates, chloroformates, reductively alkylated with amines or coupled with acids. The resulting esters were hydrolyzed to the desired acids which could further be transformed to the acylsulfonamide.


15

Method A



- 108 -


Method B



- 109 -


5

Method C




- 110 -

Method D



- 111 -

Method E



5

Example 1

10 4-({3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(phenethylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid

Step 1 - Ethyl 5-nitroindole-2-carboxylate (21.1 g) was dissolved in DMF (500 mL). To this dark brow solution, a solution of N-chlorosuccinamide was added (12 g in 125 mL DMF) over a period of 5 minutes. The reaction mixture was heated to 50 °C for 1.5 h. Reaction completion was determined by TLC. The reaction was cooled to room temperature, diluted with water (2 L), and extracted with ethyl acetate (3 X 1 L). The organic layers were combined , dried over MgSO₄, filtered and

- 112 -

5 concentrated. The result was the desired ethyl ester (96 g wet with DMF) which was carried on to the next step without further purification.

Step 2 - The ethyl ester was dissolved in methanol (400 mL) and THF (800mL). To this clear, brown solution 2 N NaOH was added (450 mL). The black
10 mixture was stirred at room temperature overnight. The reaction was not quite complete (TLC) therefore an additional 7.2 g of NaOH pellets were added. After 7 h the reaction was complete. Approximately 1 L of solvent was removed by rotary evaporation. The residue was dissolved with ethyl acetate and water and acidified with 2 N HCl to pH 2. The mixture was then extracted with ethyl acetate (3 X 1 L).
15 The combined organic layers were dried over MgSO_4 , filtered and concentrated to give the acid (24.3 g, 100%) as a brown solid.

Step 3 - The carboxylic acid (24 g) was dissolved in THF (700 mL). CDI (16.2 g) was added to this clear, amber-colored solution. The mixture was stirred at
20 room temperature for 1.5 h, during which time it became a light brown suspension. The reaction was cooled in an ice bath. Sodium borohydride (10.8 g) was added slowly. Ethanol was then added (140 mL). The evolution of gas was observed. After 2 h, TLC analysis indicated that the reaction was complete. To adjust the pH to 2, 2 N HCl was added. The reaction was then extracted with ethyl acetate (3 X 600 mL). The
25 combined organic layers were dried (MgSO_4), filtered and concentrated to yield the desired alcohol (28.5 g) as a brown solid.

Step 4 - The alcohol prepared above (25.4 g), imidazole (18.6 g), and *tert*-butyldimethylsilyl chloride (13.3 g) were dissolved in DMF (350 mL). The reaction
30 was stirred overnight and found not to be complete. Therefore imidazole (18.7 g) and *t*-butyldimethylsilyl chloride (18.6 g) were added. After 1 h, the reaction was

- 113 -

5 complete. Water (1.5 L) was added and the mixture was extracted with ethyl acetate (3 X 500 mL). The combined organic layers were evaporated to dryness to give the crude 5-*tert*-butyldimethylsilyl-protected alcohol. The crude material was dissolved in ethyl acetate, absorbed onto silica gel, and evaporated to dryness. After loading onto a silica gel column and eluting with 15% ethyl acetate in hexanes, the desired
10 protected alcohol (18.5 g, 69% from step 1) was isolated as bright yellow crystals.

Step 5 - The *tert*-Butyldimethylsilyl-protected alcohol (1.0 g) was dissolved in DMF (10 mL). The yellow solution was cooled in an ice bath. Sodium hydride (147 mg) was added. After 15 minutes, 4-(bromomethyl)benzoate (807 mg) was added to
15 the dark red solution. After 15 minutes, TLC analysis indicated that the reaction was complete. The reaction was poured into cold 1 N HCl. Water (800 mL) was added and the solution was extracted with ethyl acetate. The combined organic layers were evaporated to dryness to give the crude N-alkylated indole as an orange solid. The crude solid as absorbed onto silica gel, added to a silica gel column and eluted with
20 15% ethyl acetate in hexanes to give the desired N-alkylated indole (1.05 g, 73%) as a yellow solid.

Step 6 - N-alkylated indole (3.8 g) was dissolved in THF (50 mL) and 5% Pt/C (1.6 g) was added. Hydrogen was purged in and the reaction was stirred at room
25 temperature overnight. The reaction was filtered (celite) and concentrated. Column chromatography (35% ethyl acetate in hexane) gave the desired amine (1.7 g) as an off-white solid.

Step 7 - To a solution of the above amine (1.6 g, 3.5 mmol) in CH_2Cl_2 (10 mL)
30 and saturated sodium bicarbonate (10 mL) was added cyclopentyl carbonyl chloride (0.467 mL, 1.1 eq). The reaction was stirred 45 min, taken up in ethyl acetate (100

- 114 -

5 mL), washed with brine (3 X 20 mL), dried (MgSO_4), filtered and concentrated. Chromatography (20% ethyl acetate/hexanes) afforded the desired product (1.55 g, 82%) as a pale yellow oil.

10 Step 8 - To a solution of the above amide (0.6 g, 1.1 mmol) in dichloroethane (3 mL) at 0 °C is added dibromotriphenylphosphorane (0.5 g, 1.1 eq). The reaction is stirred to room temperature over 2h, taken up in ethyl acetate (50 mL), washed with brine (3 X 10 mL), dried (MgSO_4) and concentrated and carried immediately to the next step.

15 Step 9 - To a solution of the crude bromide prepared above (0.54 mmol) in DMF (2 mL, degassed) is added phenethyl mercaptan (0.08 g, 1.1 eq) followed by cesium carbonate (0.21 g, 1.2 eq). The reaction is stirred 1h, taken up in ethyl acetate (20 mL), washed with brine (3 X 5 mL), dried (MgSO_4), filtered and concentrated. Chromatography (25% ethyl acetate/hexanes) afforded the desired compound (0.17 g,
20 56%) as a colorless oil.

Step 10 - To a solution of the above ester in THF (1 mL) and MeOH (0.5 mL) is added NaOH (0.28 mL, 5 M, 5 eq). The reaction is stirred 4 h, acidified with sodium biphosphate, poured into ethyl acetate, washed with brine and dried (MgSO_4).
25 The title compound (0.157, 98%) was triturated from ethyl acetate with hexanes.

Each of the compounds of the following Examples 2 through 11 was prepared by a first step as illustrated in Example 1, step 9, using the appropriate thiol, followed by a second step as described in Example 1, step 10.

30

Example 2

- 115 -

5 4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(2-furylmethyl)sulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

Example 3

10 4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(4-hydroxy-6-phenyl-2-pyrimidinyl)sulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

Example 4

15 4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(4-(2-thienyl)-2-pyrimidinyl)sulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

Example 5

4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(2,4-dibromophenoxy)methyl]-1H-indol-1-yl)methyl]benzoic acid

20 Example 6

4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(cyclopentylsulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

Example 7

25 4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(propylsulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

Example 8

30 4-[(2-[(4-(tert-butyl)phenoxy)methyl]-3-chloro-5-[(cyclopentylcarbonyl)amino]-1H-indol-1-yl)methyl]benzoic acid

- 116 -

5

Example 9

4-({3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(2-quinolinylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

10 Example 10

4-[(3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(cyclopropylmethyl)sulfanyl]methyl)-1H-indol-1-yl)methyl]benzoic acid

Example 11

15 4-({2-[(benzhydrylsulfanyl)methyl]-3-chloro-5-[(cyclopentylcarbonyl)amino]-1H-indol-1-yl)methyl}benzoic acid

The compounds of the following Examples 12 through 14 were prepared by:

20 Step 1 - The material prepared in Example 1, step 6, was acylated at the 5-amino position using the protocol of Example 1, step 7, with the appropriate acylating reagent.

25 Step 2 - The title compound was prepared from the intermediate of Step 1 following the procedure described in Example 1, steps 8 through 10, using the appropriate thiol.

Example 12

30 4-({5-[(3-carboxypropanoyl)amino]-3-chloro-2-[(phenethylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

- 117 -

5

Example 13

4-[(5-[(3-carboxypropanoyl)amino]-3-chloro-2-[(3-methylbenzyl)sulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

10

Example 14

4-({2-({[4-(tert-butyl)benzyl]sulfanyl)methyl}-5-[(3-carboxypropanoyl)amino]-3-chloro-1H-indol-1-yl)methyl}benzoic acid

Example 15

15

4-({3-chloro-5-(3-furoylamino)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

The title compound was prepared as described in Example 43 using the appropriate acylating reagent.

20

The following Examples 17 through 21 were prepared as described in Example 43 using the appropriate acylating reagent.

Example 17

25

4-({3-chloro-5-[(3-(diethylamino)propanoyl)amino]-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

Example 18

30

4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(3-thienylcarbonyl)amino]-1H-indol-1-yl)methyl}benzoic acid

- 118 -

5 Example 19

4-({5-[(benzylamino)carbonyl]amino}-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

Example 20

10 4-({5-[(butylamino)carbonyl]amino}-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

Example 21

15 3-[(1-(4-carboxybenzyl)-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-5-yl)amino)carbonyl]benzoic acid

Example 22

4-[[5-(benzyloxy)-2-[(E)-2-carboxyethenyl]-3-(2-naphthoyl)-1H-indol-1-yl]methyl}benzoic acid

20 step 1

A solution of 2(tert -butyldimethylsilyloxymethyl)-5-benzyloxyindole (2.0g, 5.4mmol) in anhydrous ether (10ml) was cooled to -78°C and a solution of ethylmagnesium bromide (3.0M in ether, 4.0ml, 12.0mmol) was added dropwise. The mixture was stirred at -60°C to -65°C for 2h during which time the homogeneous solution became a yellow slurry. A solution of naphthoyl chloride (2.28g, 12.0mmol) in ether (8ml) was then added. After stirring for 2h at -60°C to -40°C the reaction was carefully quenched with saturated aqueous sodium bicarbonate and diluted with EtOAc. The combined organic layers were washed with brine, dried and concentrated. Flash chromatography (Hex/Acetone, 6/1) gave 2.2g (78%) of 17 as a yellow foam.

30

- 119 -

5 step 2

To an ice-cold (0C) solution of the intermediate above (1.0g, 1.9mmol) in DMF (10ml) was added sodium hydride (0.12g, 2.1mmol). The ice bath was removed after 10 min and the reaction was stirred at rt for 30min at which time the bromomethyl SEM ether (0.5ml, 2.8mmol) was added dropwise. The mixture was stirred at rt for 10 4h, water was added and the mixture was extracted with ethyl acetate. The organic layer was washed with brine, dried over magnesium sulfate and concentrated. Flash chromatography (Hex/Acetone, 6/1) afforded 1.22g (81%) of the desired intermediate as an off-white foam.

15 step 3

To a solution of the above (6.6g, 8.4mmol) in THF (80ml) was added tetrabutylammonium fluoride (1.0M in THF, 21ml). The reaction was stirred at rt for 2h, water was added and the mixture was extracted with EtOAc. The combined organic layers were washed with water, brine, dried and concentrated. Flash 20 chromatography (Hex/EtOAc, 4:1) gave 3.8g (67%) of the alcohol as a thick colorless oil. The alcohol was dissolved in THF (50ml) and MnO₂ (5.5g, 63.2mmol) was added. The reaction was stirred for 22h and filtered through a pad of Celite. Concentration of the filtrate gave 3.7g (96%) of the desired intermediate as a light yellow foam.

25

step 4

To an ice-cold (0°C) solution of trimethylphosphonoacetate (0.12ml, 0.7mmol) in DMF (5ml) was added sodium hydride (0.027g, 0.8mmol). After 30min a solution of the above intermediate (0.5g, 0.7mmol) in 5ml of DMF was added. The ice bath was 30 removed and the reaction was allowed to stir overnight. Water was added and the mixture was extracted with ethyl acetate. The combined organic layers were washed

- 120 -

- 5 with brine, dried over MgSO_4 and concentrated. Flash chromatography (Hex/EtOAc, 3/2) gave 0.2g of the desired intermediate (37%) as a white foam.

step 5

- 10 4-{[5-(benzyloxy)-2-[(E)-2-carboxyethenyl]-3-(2-naphthoyl)-1H-indol-1-yl]methyl}benzoic acid

- To a solution of the intermediate above (0.5g, 0.7mmol) in acetonitrile (10ml) was added aqueous 48% hydrofluoric acid (5ml). After 2h, water was added and the product was extracted with ethyl acetate. The organic layer was washed with water, brine and dried over magnesium sulfate. Concentration gave a crude solid that was
15 dissolved in THF (2ml) and MeOH (1ml) and 1N sodium hydroxide solution (2ml). After stirring overnight at rt the reaction was acidified to pH=3 with 10% HCl solution and extracted with ethyl acetate. Flash chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$, 10/1) gave the title compound (0.2g, 50%) as a white solid.

- 20 **Example 23**

4-([3-acetyl-5-(benzyloxy)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl]methyl)benzoic acid

- Step 1: Ethyl 5-benzyloxy-2-indolcarboxylate (30 g, 102 mmol) is dissolved in 250 mL of THF and cooled to 0° C and Lithium Aluminum Hydride (LAH) (255
25 mL of a 1.0 M solution in THF) is added via addition funnel over 40 minutes. The reaction was stirred a further 2 hours at 0° C and then worked up by the addition of 4N NaOH (190 mL). The resulting salts are filtered and washed with ethyl acetate (3X400 mL), the filtrates are combined and dried over MgSO_4 and concentrated to yield 24.8 g.

- 30

- 121 -

5 Step 2: The crude indole alcohol prepared in step 1 (8.3 g, 32.6 mmol) is dissolved in DMF (10.5 mL). To this solution is added imidazole (5.5g, 81.5 mmol) and t-butyldimethylsilyl chloride (5.4g, 35.8 mmol). The mixture is stirred at room temperature overnight. The reaction is poured into water and extracted with ethyl acetate (3X). Organic layers are dried over magnesium sulfate and concentrated. The
10 crude material is purified on a silica gel column using 19:1 Hexane:Ethyl acetate to give pure product (11.9g, 31 mmol, 94 % yield, TLC \approx 0.8 Rf in Toluene:Ethyl acetate 2:1)

Step 3: A solution of the silyl protected indole prepared in step 2 (2 g, 6.56
15 mmol) in ether (20 mL) was added dropwise to ethyl magnesium bromide (2.4 mL of a 3M solution in ether, 7.2 mmol) in ether (10 ml), the latter maintained at -78°C . The reaction was stirred at -60°C for 2 hr. Next acetylchloride (0.51 mL, 7.2 mmol) in ether (4 mL) was added slowly. The reaction maintained between -50°C and -60°C for another 2 hrs. The reaction was then quenched with saturated sodium bicarbonate.
20 Extracted with ethyl acetate (3X). Organic layers were dried over magnesium sulfate and concentrated. The crude material was purified on a silica gel column using 19:1 Hexane:Ethyl acetate to give pure product (1.2 g, 50%).

Step 4: To the indole (1.2 g, 2.9 mmol, prepared in step 3 above) in 10.5 ml of
25 DMF, sodium hydride (0.13g, 60% oil dispersion, 3.23 mmol) is added at room temperature. The reaction is stirred for 30 minutes. Methyl (4-bromomethyl) benzoate (0.81 g, 3.53 mmol) is added at this time and the reaction stirred overnight. On completion of the reaction (monitored by TLC) it is quenched with water, extracted with ethyl acetate (3X). Organic layers are dried over magnesium sulfate,
30 concentrated and used for the next step.

- 122 -

5 Step 5: A mixture of silyl protected indole prepared in step 4 above (0.65 g, 1.2 mmol) and tetra-butyl ammonium fluoride (2.9 mL of a 1M solution in tetrahydrofuran, 2.9 mmol) in tetrahydrofuran (6 mL) were stirred at room temperature for one hour. At this time the reaction was diluted with ethyl acetate and water, extracted with ethyl acetate (3X), dried over magnesium sulfate and concentrated. The crude material was purified on silica gel using 1:1 Hexane:Ethyl acetate to yield pure alcohol (0.47 g, 91 %).

15 Step 6: The indole alcohol (0.3 g, 0.68 mmol), carbon tetrabromide (0.27 g, 0.81 mmol) and 1,3-bis(diphenylphosphino)propane (0.21 g, 0.51 mmol) were taken up in dichloromethane (8.4 mL) and stirred for 16 hours at which time the reaction was diluted with dichloromethane and half saturated ammonium chloride. The aqueous layer was extracted with ethyl acetate (3X) dried over magnesium sulfate and concentrated. The crude material was purified on silica gel using 2:1 Hexane:Ethyl acetate to yield pure alcohol (0.27 g, 78 %).

20 Step 7: The indole bromide prepared in step 6 (0.1 g, 0.2 mmol) was dissolved in dimethylformamide (0.4 mL, degassing the solvent is strongly recommended) cesium carbonate (0.2 g, 0.6 mmol) was added and then ethyl 2 naphthalenethiol (0.034g, 0.22 mmol) was added and the mixture was stirred for 1 day, then the reaction was poured into 1/2 saturated ammonium chloride and extracted with ethyl acetate (3X), dried, concentrated and chromatographed (Hexane:Ethyl acetate 3:1) to yield 0.05 g (57%) of pure product.

30 Step 8: The ester (0.2 g, 0.34 mmol) prepared in step 7 above is dissolved in 4.0 mL of 1/1 THF/ methanol and then 1N sodium hydroxide (2.5 mL) is added and the resulting mixture is stirred for 16 hours at RT, workup gave crude product that is

- 123 -

5 purified via chromatography (1:1 Hexane:Ethyl acetate with 1% acetic acid) to yield (0.17 g, 85%) of solid.

Example 24

10 **4-[[5-(benzyloxy)-2-[(2-naphthylsulfanyl)methyl]-3-(2,2,2-trifluoroacetyl)-1H-indol-1-yl]methyl}benzoic acid**

Step 1: This intermediate was prepared from the indole, prepared in step 2 of Example 23 and trifluoroacetic anhydride, according to the procedure described in step 3 of Example 23.

15 Step 2: This intermediate was prepared according to the procedure described in step 4 of Example 23, but using the indole derivative prepared in step 1 above and methyl (4-bromomethyl) benzoate.

20 Step 3: To a solution of the indole alcohol prepared in step 2 (0.1 g, 0.2 mmol) and triethylamine (0.04 mL, 0.3 mmol) in dichloromethane (0.4 mL), methanesulphonyl chloride (0.02 mL, 0.24 mmol) is added dropwise at 0°C. The reaction is stirred for 1.5h and then the dichloromethane removed. To the residue in 0.4 mL of DMF at 0°C 2-naphthalene thiol (0.034 g, 0.22 mmol) is added. Next CsCO₃ (0.96 g, 0.3 mmol) is added and the reaction mixture stirred at room temperature overnight, then the
25 reaction was poured into brine and extracted with ethyl acetate (3X), dried, concentrated and chromatographed (Hexane:Ethyl acetate 3:1) to yield 0.064 g (50%) of pure product.

30 Step 4: The title compound was prepared from ester, prepared in step 3 above, according to the procedure described in step 8 of Example 23.

- 124 -

5 Example 25

4-({5-[(4-aminobutanoyl)amino]-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

step 1: The t-butylsilyl protected alcohol (25g) from step 6 of Example 1, was dissolved in THF(200mL) and then tetrabutylammonium fluoride (125mL of a 1.0 M solution) was added and the mixture was stirred for 10 minutes at room temperature and then the reaction was diluted with water and the THF was concentrated, ethyl acetate was added and the layers were separated, the aqueous layer was extracted three times with ethyl acetate, the combined organic layers were dried and concentrated to yield the desired alcohol (21.5 g).

Step2: The alcohol (13.1g) from step 1 was suspended in dichloromethane (450 mL), cooled to 0°C and triethylamine(10 mL) and methanesulfonyl chloride (4.0 mL) were added and the resulting mixture was left to warm to room temperature overnight at which time it was diluted with saturated sodium bicarbonate was added and the reaction was diluted with dichloromethane, the layers were separated, the aqueous layer was extracted three times with dichloromethane, the combined organic layers were dried, concentrated to yield the desired chloride (13.5g).

Step 3: To the chloride (13.5g) generated in step 2 was added DMF (150 mL), cesium carbonate (33.5 g) and then the solution was degassed by bubbling argon through the solution for 20 minutes and then 2-naphthalene thiol was added and the reaction was stirred for 20 minutes at room temperature, then water was added as was ethyl acetate, the layers were separated and the combined organic layers were concentrated to a slurry which was stirred overnight and then the slurry was filtered and the solid was triturated with 40% ethyl acetate in hexane to deliver the desired disulfide (12.2 g) in 69% yield.

- 125 -

5

Step 4: The product from step three (11.25 g) was dissolved in THF (500 mL), methanol (500mL) and then copper II acetate (19.2 g) suspended in water (300 mL) was added as was more THF (100mL) and then sodium borohydride (11.2 g) was added portionwise. After 2.5 hours of stirring at room temperature the foamy black solution was diluted with saturated sodium bicarbonate and the layers were separated, the aqueous layer was extracted three times with ethyl acetate, the combined organic layers were dried and concentrated and chromatographed to yield the desired amine (9.0 g) in 85% yield.

10

15

step 5: The amine from step 4 above was coupled to fmoc protected 4-aminobutyric acid following the procedure outlined in step 1, Example 43, followed by trituration with dichloromethane delivered the amide in 43% yield.

20

Step 6 : The amide (1.0 eq) from step 5 was dissolved in methanol (5 mg/ml) and piperidine (0.024 ml/mg) and then the reaction was stirred at room temperature for two hours, concentrated and chromatographed to yield the desired product in quantitative yield.

25

Step 7: The amino ester from step 6 was hydrolyzed using the conditions outlined for step 2 of Example 43 to deliver the title compound in 54%.

Example 26

4-({3-chloro-5-[(cyclopentylcarbonyl)amino]-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

30

step1: The amine (1.0 eq) from step 4 of Example 25 was dissolved in CH₂Cl₂ (0.3M) and then an equivalent amount of saturated sodium bicarbonate was added and then

- 126 -

- 5 the appropriate acid chloride (1.2 eq) was added. The biphasic reaction mixture was vigorously stirred until TLC analysis indicated that the reaction was complete (generally a few hours) and then the reaction was diluted with dichloromethane and water, the layers were separated, the aqueous layer was extracted three times with dichloromethane, the combined organic layers were dried, concentrated and
10 chromatographed, or used crude, to yield the desired amide in 50% yield.
- Step 2: The ester from the previous step was dissolved in THF/MeOH (3:1) and then 1N NaOH (3.0eq) was added and the reaction was stirred for until TLC analysis showed that the reaction was complete. The reaction was then concentrated, diluted with water, acidified to pH 2 with conc HCL, extracted with ethyl acetate 3X, the
15 combined organics were dried over magnesium sulfate concentrated and purified via chromatography to yield the desired acid in 69% yield.

Example 27

- 20 **4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(2-quinoxalinylicarbonyl)amino]-1H-indol-1-yl)methyl}benzoic acid**

Step1: The amine from step 4 of Example 25 was treated with the appropriate acid chloride according to the general procedure for Example 71, step 1, to deliver the amide in 76% yield.

- 25 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 53% yield.

Example 28

- 30 **4-({3-chloro-5-[(2,2-dimethylpropanoyl)amino]-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid**

- 127 -

5 Step1: The amine from step 4 of Example 25 was treated with the appropriate acid chloride according to the general procedure for Example 71, step 1, to deliver the amide in •100% yield.

10 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 79% yield.

Example 29

4-({5-}[(benzyloxy)carbonyl]amino)-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid

15 step1: The amine from step 4 of Example 25 was treated with the appropriate acid chloride according to the general procedure for Example 71, step 1, to deliver the amide in •96% yield.

20 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid.

Example 30

4-({3-chloro-5-}[(cyclopentyloxy)carbonyl]amino)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid

25 Step1: The amine from step 4 of Example 25 was treated with acetic anhydride according to the general procedure for Example 71, step 1, to deliver the amide in 92% yield.

30 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid.

5 **Example 31**

4-({5-(acetylamino)-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid

Step1: The amine from step 4 of Example 25 was treated with acetic anhydride according to the general procedure for Example 71, step 1, to deliver the amide in 77% yield.

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 29%.

15 **Example 32**

4-({5-[(butylamino)carbonylamino]-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid

Step1: THF (0.12M) was added to the amine (1.0 eq) generated in step 4, the reaction mixture was cooled to 0°C and then butylisocyanate (1.1 eq) was added and the mixture was warmed to room temperature overnight and the reaction was diluted with 1/2 saturated ammonium chloride, the layers were separated, the aqueous layer was extracted three times with ethyl acetate, the combined organic layers were dried and concentrated and purified via chromatography to yield the desired urea in 57% yield.

25

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 41%.

Example 33

30 **4-({5-[(butylamino)carbonylamino]-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl)benzoic acid**

- 129 -

5 Step 1: According to the procedure for Example 32, step 1, the amine was treated with benzyl isocyanate to deliver the title compound in 16% yield.

 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 100%.

10

Example 34

4-({3-chloro-5-[(morpholinocarbonyl)amino]-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

 Step 1: The amine (1.0 eq) generated above in step 4 was weighed into a flask
15 along with 4-dimethylaminopyridine (1.5eq) and then they were taken up in dichloroethane (0.08 M) and then 4-morpholinecarbonyl chloride (1.5eq) was added and the reaction was stirred overnight at room temperature and then heated to 40°C for 4 hours and then worked up by the addition of ethyl acetate and 1/2 saturated ammonium chloride, the layers were separated, the aqueous layer was extracted three
20 times with ethyl acetate, the combined organic layers were dried and concentrated and purified via chromatography to yield the desired urea in •100% yield.

 Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 79%.

25

Example 35

4-({5-(benzylamino)-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

 Step 1: The amine (1.0eq) from step 1 was dissolved in dichloroethane and
30 then benzaldehyde (1.0 eq) was added followed by acetic acid (1.0 mL/1mmol) and the reaction was stirred for 20 minutes and then the sodium triacetoxyborohydride

- 130 -

5 (1.3 eq) was added and the reaction was stirred overnight at room temperature, quenched by the addition of aqueous diethanolamine and dichloromethane the layers were separated, the aqueous layer was extracted three times with dichloromethane , the combined organic layers were dried and concentrated and purified via chromatography to yield the desired urea in 74% yield.

10

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 49%.

Example 36

15 **4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(3-phenoxybenzyl)amino]-1H-indol-1-yl)methyl}benzoic acid**

Step 1: According to the procedure for step 1 of Example 35 the amine from step 4 of Example 25 was treated with the appropriate aldehyde to yield the desired secondary amine in 38% yield.

20

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 87%.

Example 37

25 **4-({3-chloro-5-[(cyclopentylcarbonyl)(methyl)amino]-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid**

Step 1: To the ester generated in step 1 of the synthesis of Example 27 was added DMF (0.05M), the reaction was cooled to 0°C and then sodium hydride (10 eq) was added and the mixture was stirred for 30 minutes, methyl iodide (10eq) was then added and the resulting mixture was stirred overnight at room temperature and then diluted with ethyl acetate and 1/2 saturated ammonium chloride, the layers were

30

- 131 -

5 separated, the aqueous layer was extracted three times with ethyl acetate, the combined organic layers were dried and concentrated and purified via chromatography to yield the desired methylated amide in 56% yield.

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for
10 step 2 of Example 26 to yield the desired acid in 57%.

Example 38

4-({5-[acetyl(benzyl)amino]-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl}methyl)benzoic acid

15 Step1: The amide synthesized in step 1 of Example 31 was benzylated according to the procedure in step 1 for Example 37 to yield the tertiary amide in 90% yield.

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for
20 step 2 of Example 26 to yield the desired acid in 41%.

Example 39

4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(tetrahydro-3-furanylcarbonyl)amino]-1H-indol-1-yl}methyl)benzoic acid

25 Step 1: To the indole amine (1.0 eq) was added the acid (1.2 eq) the dimethylaminopyridine (10 mol %), 1-(3dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (1.5 eq) and then DMF(0.3M) and the reaction was stirred under nitrogen for 24 hours at room temperature at which time it was poured into 1/2 saturated ammonium chloride solution and ethyl acetate and then the layers were
30 separated and the aqueous layer was extracted 3 times, the combined organic layers

- 132 -

5 were washed with water 2X, dried, concentrated and chromatographed to yield 55% of the title compound.

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 55%.

10

Example 40

4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(3-thienylcarbonyl)amino]-1H-indol-1-yl)methyl}benzoic acid

step 1: According to the procedure for step 1 of Example 39 the amine from step 4 of Example 25 was treated with the requisite acid to yield the amide in •100% yield.

15

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 21%.

20

Example 41

4-({3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(1-adamantylcarbonyl)amino]-1H-indol-1-yl)methyl}benzoic acid

Step 1: According to the procedure for step 1 of Example 39 the amine from step 4 of Example 25 was treated with the requisite acid to yield the amide in •100% yield.

25

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 35%.

30

Example 42

- 133 -

5 3-[(1-(4-carboxybenzyl)-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-5-yl)amino]carbonylbenzoic acid

Step 1: According to the procedure for step 1 of Example 39 the amine from step 4 of Example 25 was treated with the requisite acid to yield the amide in •100% yield.

10

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 20%.

Example 43

15 4-[(3-chloro-2-[(2-naphthylsulfanyl)methyl]-5-[(3-phenylpropanoyl)amino]-1H-indol-1-yl)methyl]benzoic acid

Step 1: According to the procedure for step 1 of Example 39 the amine from step 4 of Example 25 was treated with the requisite acid to yield the amide in •100% yield.

20

Step 2: The ester from step 1 was hydrolyzed under the conditions outlined for step 2 of Example 26 to yield the desired acid in 32%.

Example 44

25 4-[(5-amino-3-chloro-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl]benzoic acid

step 1: The amine generated in step 4 was hydrolyzed according to the procedure for step 2 of Example 26 to yield 79% yield.

30 Example 45

- 134 -

5 N-{3-chloro-1-(4-[(methylsulfonyl)amino]carbonyl)benzyl)-2-[(2-naphthylsulfonyl)methyl]-1H-indol-5-yl}cyclopentanecarboxamide

Step 1: To Example 26 (1.0 eq), EDCI (1.5 eq), DMAP (1.0 eq), methane sulfonamide (1.0 eq) in a flask under nitrogen was added DMF (0.08M) and the reaction was stirred overnight at room temperature and then worked up by the addition of 1/2 saturated ammonium chloride solution and ethyl acetate and then the layers were separated and the aqueous layer was extracted 3 times, the combined organic layers were washed with water 2X, dried, concentrated and chromatographed to yield 27% of the title compound.

15 Example 46

N-{3-chloro-2-[(2-naphthylsulfonyl)methyl]-1-[4-[(4-nitrophenyl)sulfonyl]amino}carbonyl)benzyl]-1H-indol-5-yl}cyclopentanecarboxamide

Step 1: To Example 26 was added the appropriate sulfonamide under the conditions outlined for step 1, Example 45, to yield the desired acylsulfonamide in 43% yield.

Example 47

N-{3-chloro-1-[4-[(2-methylphenyl)sulfonyl]amino}carbonyl)benzyl]-2-[(2-naphthylsulfonyl)methyl]-1H-indol-5-yl}cyclopentanecarboxamide

Step 1: To Example 26 was added the appropriate sulfonamide under the conditions outlined for step 1, Example 45, to yield the desired acylsulfonamide in 40% yield.

Example 48

30 N-[3-chloro-2-[(2-naphthylsulfonyl)methyl]-1-(4-[(phenylsulfonyl)amino]carbonyl)benzyl]-1H-indol-5-yl}cyclopentanecarboxamide

- 135 -

5 Step 1: To Example 26 was added the appropriate sulfonamide under the conditions outlined for step 1, Example 45, to yield the desired acylsulfonamide in 40% yield.

Example 49

10 **N-{3-chloro-2-[(2-naphthylsulfanyl)methyl]-1-[4-({[(trifluoromethyl)sulfonyl]amino}carbonyl)benzyl]-1H-indol-5-yl}cyclopentanecarboxamide**

 Step 1: To Example 26 was added the appropriate sulfonamide under the conditions outlined for step 1, Example 45, to yield the desired acylsulfonamide in 67% yield.

15 **Example 50**

4-[5-[(cyclopentylcarbonyl)amino]-2-[(2-naphthyloxy)methyl]-3-(1-pyrrolidinylcarbonyl)-1H-indol-1-yl]butanoic acid

Step 1

20 The Ethyl 5-nitroindole-2-carboxylate (1 eq) was dissolved in THF/MeOH/H₂O (3:1:1 0.21 M) followed by addition of LiOH·H₂O (1.38 eq), stirred at 25°C, overnight. Reaction mixture was then acidified to pH=1 with 1N HCl solution and extracted with ethyl acetate. Workup with water, brine, drying and concentration afforded the crude product in 99% yield.

25 **Step 2**

 The crude acid (1 eq) from step 1 was dissolved in THF (0.14 M) and to this was added carbonyl diimidazole (1 eq), stirred at 25°C for 1.5h. Reaction mixture was then cooled to 0°C. Sodium borohydride (2.86 eq) was added in several portions
30 followed by the addition of EtOH (0.71 M), stirred at 25°C, overnight. Reaction mixture at this point was acidified with 2N HCl to pH=2 and extracted with ethyl

- 136 -

- 5 acetate. Workup with water, brine, drying and concentration afforded the crude product in 95% yield.

Step 3

- 10 In an oven dried flask was added the crude indole alcohol (1 eq) from step 2 followed by anhydrous DMF (0.135 M). To this was then added imidazole (1.3 eq) and TBSCl (1.2 eq), stirred at 25°C for 1 h. Workup with ethyl acetate/water followed by chromatographic purification afforded the desired product in 70% yield.

Step 4

- 15 The silyl protected indole (1 eq) from step 3 was dissolved in anhydrous DMF (0.13 M) in an oven dried flask. To this was added NaH (60% dispersion in mineral oil, 1.2 eq) and stirred at 25°C for 1 h after which ethyl 4-bromobutyrate (1.2 eq) and KI (1.2 eq) were added. Reaction mixture was then heated at 60°C for 2 h. Workup with ethyl acetate/water followed by chromatographic purification afforded the desired
20 product in 93% yield.

Step 5

- Dissolved the alkylated indole (1 eq) from step 4 in THF (0.05 M) and to this was added TBAF (1.0 M/THF, 1.1eq) dropwise at 25°C, stirred for 1 h. Workup with
25 ethyl acetate/water followed by chromatographic purification afforded the desired product quantitatively.

Step 6

- 30 Dissolved the indole alcohol (1 eq) from step 5 in anhydrous DMF (0.16 M) followed by the addition of POCl₃ (10 eq) dropwise at 0°C. Heated at 80°C, overnight. Workup with ethyl acetate/water and washing the organic layer with 1N NaOH,

- 137 -

- 5 water, brine followed by concentration and chromatographic purification afforded the desired product in 64% yield.

Step 7

- 10 A mixture of the chloro aldehyde derivative (1 eq) from step 6, 325 mesh K_2CO_3 (2.4 eq), 2-naphthol (1.2 eq) and KI (1.2 eq) was suspended in anhydrous acetonitrile (0.2 M) and heated at 70°C for 1.5 h. Workup with ethyl acetate/water followed by chromatographic purification afforded the desired product in 65% yield.

Step 8

- 15 Dissolved the naphthyloxy indole derivative (1 eq) from step 7 in anhydrous THF (0.023 M) and to this was added 5% Pt/C (40wt%) under nitrogen and hydrogenated with a H_2 balloon for 2.5 h. Reaction mixture was then filtered through Celite and concentrated to afford the crude product in 96% yield.

20 **Step 9**

- The amino indole derivative (1 eq) from step 8 was dissolved in anhydrous CH_2Cl_2 (0.07 M) followed by dropwise addition of Et_3N (1.4 eq) and cyclopentanecarbonyl chloride (1.2 eq) at 0°C. Stirred at 25°C for 0.5 h after which the reaction mixture was quenched with saturated $NaHCO_3$ solution and stirred overnight. Organic layer was separated and washed with brine, dried. Product was obtained in 66% yield after recrystallization from 30% ethyl acetate/hexane.

Step 10

- 30 The indole from step 9 (1 eq) was weighed into a flask along with NaH_2PO_4 (12 eq), t-butyl alcohol (0.13 M), water (0.13 M), 2-methyl-2-butene (46 eq) and to this mixture was added NaO_2Cl (12 eq) at 25°C. Reaction mixture was then heated at

- 138 -

5 65°C, overnight. Workup with ethyl acetate/water followed by trituration with CH₂Cl₂/hexane (4:6) at 0°C for 1 h afforded the desired product in 63% yield.

Step 11

10 The acid (1 eq) from step 10 was weighed into a flask and to this was added EDCI (3 eq), DMAP (1.2 eq), pyrrolidine (1.2 eq) followed by anhydrous THF (0.018 M) and the reaction mixture was then refluxed for 18 h. Worked up with ethyl acetate/water followed by washing the organic layer with 1N HCl, saturated bicarbonate and brine. Recrystallization from ethyl acetate/hexane (3:7) afforded the desired product in 89% yield.

15

Step 12

Dissolved the amide (1 eq) from step 11 in THF/MeOH/water (3:1:1, 0.025 M) and to this was added LiOH·H₂O (1.2 eq), stirred at 25°C, overnight. Workup with ethyl acetate/water followed by recrystallization from CH₂Cl₂/hexane (1:1) afforded the desired product in 98% yield.

20

Example 51

4-{5-[(cyclopentylcarbonyl)amino]-3-(morpholinocarbonyl)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoic acid

25 Step 1

Following step 11 of Example 50 and using morpholine yielded 87% of the desired product after recrystallization.

Step 2

30 Following step 12 above (Example 50) and using the corresponding morpholino amide afforded the desired product in 96% yield after recrystallization.

5 Example 52

N-[2-[(2-naphthyloxy)methyl]-1-(4-oxo-4-
{[(trifluoromethyl)sulfonyl]amino}butyl)-3-(1-pyrrolidinylcarbonyl)-1H-indol-5-
yl]cyclopentanecarboxamide

10 The acid Example 50 (1 eq) was weighed into a flask and to this was added
EDCI (3 eq), DMAP (1.2 eq), trifluoromethanesulfonamide (1.2 eq) followed by
anhydrous THF (0.04 M) and the reaction mixture was stirred at 25°C, overnight.
Worked up with ethyl acetate/water followed by washing the organic layer with 1N
HCl, saturated bicarbonate and brine. Trituration of crude product with
CH₂Cl₂/hexane (1:2) at 0°C for 1 h afforded the desired product in 96% yield.

15

Example 53

N-[3-(morpholinocarbonyl)-2-[(2-naphthyloxy)methyl]-1-(4-oxo-4-[(trifluoro-
methyl)sulfonyl]amino}butyl)-1H-indol-5-yl]cyclopentanecarboxamide

20 Following step 1 above (Example 52) and using the corresponding acid
Example 51 (step 4, scheme-4) afforded the desired product in 96% yield.

Example 53A

4-{5-[(cyclopentylcarbonyl)amino]-3-formyl-2-[(2-naphthyloxy)methyl]-1H-
indol-1-yl}butanoic acid

25 Step 1

Dissolved the indole (1 eq) from step 9 Example 50 in THF/MeOH/H₂O
(3:1:1, 0.025 M) and to this was added LiOH.H₂O (1.2 eq), stirred at 25°C, for 4 h.
Workup with ethyl acetate/1N HCl followed by trituration with CH₂Cl₂/hexane
afforded the desired product in 74% yield.

30

- 140 -

5 Example 53 B

N-[3-formyl-2-[(2-naphthyloxy)methyl]-1-(4-oxo-4-
{[(trifluoromethyl)sulfonyl]amino}butyl)-1H-indol-5-
yl]cyclopentanecarboxamide

Step 1

10 The acid (1 eq) from step 1 of **Example 53A** was weighed into a flask and to
this was added EDCI (1.35 eq), DMAP (1.1 eq), trifluoromethanesulfonamide (1.05
eq) followed by anhydrous THF (0.026 M) and the reaction mixture was stirred at
25°C, 3 h. Worked up with ethyl acetate/water followed by washing the organic layer
15 with 0.05N HCl, saturated bicarbonate and brine. Chromatographic purification then
afforded the desired product in 94% yield.

Example 54

5-[(cyclopentylcarbonyl)amino]-2-[(2-naphthyloxy)methyl]-1-(4-oxo-4-
{[(trifluoromethyl)sulfonyl]amino}butyl)-1H-indole-3-carboxylic acid

20 Step 1

The product of **Step 1, Example 53B** (1 eq) was weighed into a flask along
with NaH₂PO₄ (12 eq), t-butyl alcohol (0.12 M), water (0.12 M), 2-methyl-2-butene
(50 eq) and to this mixture was added NaO₂Cl (11.8 eq) at 25°C. Reaction mixture
was then heated at 60°C, 3 h and left overnight at 25°C. Workup with ethyl
25 acetate/water followed by chromatographic purification and triturations with
CH₂Cl₂/hexane (1:1) afforded the desired product in 57% yield.

Example 55

3-({4-[5-[(cyclopentylcarbonyl)amino]-2-[(2-naphthyloxy)methyl]-3-(1-
30 pyrrolidinylcarbonyl)-1H-indol-1-yl]butanoyl}amino)benzoic acid

- 141 -

5 **Step 1**

 The compound of Example 50 (1 eq) was weighed into a flask and to this was added EDCI (3 eq), DMAP (1.2eq), methyl 3-aminobenzoate (1.2 eq) followed by anhydrous THF (0.04M) and the reaction mixture was then stirred at 25°C for 2 d. Worked up with ethyl acetate/water followed by washing the organic layer with 1N HCl, saturated bicarbonate and brine. Recrystallization from ethyl acetate/hexane afforded the desired product in 88% yield.

10

Step 2

 Dissolved the ester (1 eq) from step 1 in THF/MeOH/water (3:1:1, 0.024 M) and to this was added LiOH·H₂O (1.2 eq), stirred at 25°C, overnight at this point 1.2 eq of LiOH·H₂O was added and stirred for 2 h. Workup with ethyl acetate/1N HCl followed by trituration with CH₂Cl₂/hexane (1:1) at 0°C for 1h afforded the desired product in 92% yield.

15

20 **Example 56**

3-[(4-{5-[(cyclopentylcarbonyl)amino]-3-(morpholinocarbonyl)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)amino]benzoic acid

Step 1

 Following step 1 of Example 55 and using the product of Example 51 (see scheme-4 for synthesis) afforded the desired product in 85% yield after chromatographic purification.

25

Step 2

 Following step 2 of Example 55 and using the corresponding morpholino from step 1 afforded the desired product in 91% yield.

30

Example 57

- 142 -

5 N-[2-[(2-naphthyloxy)methyl]-1-[4-oxo-4-[3-
([(trifluoromethyl)sulfonyl]amino)carbonyl]anilino]butyl]-3-(1-
pyrrolidinylcarbonyl)-1H-indol-5-yl]cyclopentanecarboxamide
Step 1

10 The compound of Example 55 (1 eq) was weighed into a flask and to this was
added EDCI (3 eq), DMAP (1.2 eq), trifluoromethanesulfonamide (1.2 eq) followed
by anhydrous THF (0.04 M) and the reaction mixture was then stirred at 25°C,
overnight. Worked up with ethyl acetate/water followed by washing the organic layer
with 1N HCl, saturated bicarbonate and brine. Trituration with CH₂Cl₂/hexane (8:2)
afforded the desired product in 84% yield.

15

Example 58

N-(3-(morpholinocarbonyl)-2-[(2-naphthyloxy)methyl]-1-[4-oxo-4-[3-
([(trifluoromethyl)sulfonyl]amino)carbonyl]anilino]butyl)-1H-indol-5-
yl)cyclopentanecarboxamide

20 Following step 1 of Example 57 and using Example 56 afforded the desired
product in 84% yield.

Example 59

2-(4-[5-(benzyloxy)-3-(1-naphthoyl)-1H-indol-1-yl]methyl}phenyl)acetic acid

25 Step 1: A solution of MeMgBr in butyl ether (1 M, 1.2 eq) was ice cooled. 4-
Benzyloxy indole (1 eq) in CH₂Cl₂ (0.5 M) was added and the reaction mixture was
allowed to warm to 25°C. After the addition of 1-naphthoyl chloride (1 eq) in CH₂Cl₂
(1 M) the reaction was heated to reflux for 3 h. Quenching with aqueous NH₄Cl and
extraction with CHCl₃ provided crude ketone, which was purified by recrystallization
30 from hexane/CHCl₃/MeOH (53 % yield).

- 143 -

5 Step 2: An ice cooled solution of the ketone (1 eq) from step 1 in DMF (0.2 M) was treated with NaH (60 % in mineral oil, 2.5 eq). 4-bromophenyl acetic acid (1.1 eq) in DMF (0.4 M) was added after 15 minutes and the resulting mixture was stirred overnight at 25°C. The reaction was quenched with 1N HCl and extracted with EtOAc. The organic extracts were dried and concentrated. The desired product
10 was obtained in 68 % yield after purification by chromatography and recrystallization from hexane/EtOAc.

Example 60

2-(4-([5-(benzyloxy)-3-(2-naphthoyl)-1H-indol-1-yl]methyl)phenyl)acetic acid

15 Step 1: Following step 1 of Example 59 using the appropriate acyl chloride yielded 42 % of the desired ketone after recrystallization from hexane/CHCl₃.

 Step 2: An analogous procedure to step 2 of Example 59 yielded 35 % of the title compound after chromatographic purification and recrystallization from acetone/pentane.

20 **Example 61**

2-[4-([5-(benzyloxy)-3-[3,5-bis(trifluoromethyl)benzoyl]-1H-indol-1-yl]methyl)phenyl]acetic acid

25 Step 1: Following step 1 of Example 59 using the appropriate acyl chloride yielded 30 % of the desired ketone after recrystallization hexane/CH₂Cl₂/EtOAc.

 Step 2: An analogous procedure to step 2 of Example 59 yielded 73 % of the title compound after chromatographic purification and recrystallization from CHCl₃/MeOH.

30 **Example 62**

- 144 -

5 4-({3-benzoyl-5-(benzyloxy)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

Step 1 The starting ethyl 5-benzyloxyindole-2-carboxylate (Scheme 21, step 1) was treated with LAH (1.3 eq) in THF (0.27 M) at 0°C under nitrogen for 1 h. Workup with NaOH and water followed by concentration afforded crude product (100%).

10

Step 2 The crude alcohol from step 1 was dissolved in DMF (0.38 M), and treated with t-butyldimethylsilyl chloride (1.16 eq) and imidazole (1.26 eq) at 25°C for 1 d. Workup and chromatographic purification afforded the pure product (93%).

15 Step 3 The silyl ether from step 2 was dissolved in methylene chloride (0.26 M), and treated with BOC anhydride (1.24 eq), triethylamine (1.53 eq) and DMAP (0.21 eq) at 25 °C for 3 d. Workup and chromatographic purification afforded the pure product (99%).

20 Step 4 The N-BOC silyl ether from step 3 was treated with acetic acid/ water/ THF (3:1:1) (0.04 M) at 25°C for 1 d. Workup and chromatographic purification afforded the pure product (100%).

Steps 5 The alcohol from step 4 was dissolved in methylene chloride (0.2 M), and under nitrogen at -40°C treated with triethylamine (1.33 eq), and mesyl chloride (1.23 eq) for 1 h. In a separate dry flask was weighed naphthalene-2-thiol (1.31 eq), and THF (1 M) was added, followed by lithium hexamethyldisilazide (1N in THF, 1 eq) and this mixture was stirred at 25°C for 30 min. The resulting solution was then added dropwise, over 30 minutes, to the above mesylate solution, at -40°C. The reaction mixture was allowed to warm to 25°C, and stirred there for 4.5 h. Workup and chromatographic purification afforded the BOC thioether.

25

30

- 145 -

5

Step 6 The purified BOC thioether from step 5 was heated under nitrogen at 160-170°C for 1.25 h, and recrystallized from ethyl acetate and hexanes to afford the free indole thioether in 64% yield.

10 Step 7: The product of step 6 (1 eq) in CH_2Cl_2 (0.125 M) was ice cooled. A solution of MeMgBr (1.2 eq) in butyl ether (1 M) was added and the resulting mixture was stirred for 30 minutes. After warming up to 25°C, benzoyl chloride was added dropwise. The reaction was heated to reflux for 3 h, then stirred at 25°C overnight. After quenching with NH_4Cl , the mixture was extracted with CH_2Cl_2 . Organic
15 extracts were washed with brine, dried and concentrated. The desired ketone was obtained in 55 % yield after chromatographic purification.

Step 8: A solution of the product from step 1 (1 eq) in dry DMF (0.1 M) was treated with NaH (60 % in mineral oil, 1.05 eq). Methyl 4-bromomethylbenzoate (1.2 eq)
20 was added after 1 h at 25°C and the resulting mixture was stirred overnight. EtOAc /water work up produced the desired crude material which was purified by chromatography (56 % yield).

Step 9: The material from step 2 (1 eq) was hydrolyzed by the action of $\text{LiOH}\cdot\text{H}_2\text{O}$
25 (1.2 eq) in $\text{THF}/\text{MeOH}/\text{water}$ (3/1/1, 0.07 M). After stirring at 25°C overnight, the reaction mixture was quenched with AcOH and solvent was evaporated. EtOAc /water work up and chromatographic purification afforded the title compound in 78 % yield.

30 **Example 63**

- 146 -

5 4-({5-(benzyloxy)-3-isobutyryl-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoic acid

Step 1: Following step 7 of Example 62 using isopropyl chloride yielded the desired ketone after chromatography.

10 Step 2: An analogous procedure to step 8 of example 62 yielded 50 % of the N-alkylated material after chromatographic purification.

Step 3: Following step 9 of example 62 the methyl ester was hydrolyzed to the title compound in 67 % yield after chromatography.

Example 64

15 2-{3-acetyl-5-(benzyloxy)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl}acetic acid

Step 2: The product of step 1 of Example 80 was alkylated with methyl bromoacetate following an analogous procedure to step 8 of example 62 to yield 65 % of the desired compound after recrystallization from EtOAc.

20 Step 3: Following step 9 of example 62 above the methyl ester was hydrolyzed to the title compound in 84 % yield after chromatography.

Example 65

25 2-{5-(benzyloxy)-3-isobutyryl-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl}acetic acid

Step 2: An analogous procedure to step 2 of Example 63 started from methyl bromo acetate and the isopropyl ketone described in step 1 of Example 63 (step 1, see above). This reaction yielded 66 % of the N-alkylated material after chromatographic purification.

30 Step 3: Following step 3 of Example 63 the methyl ester was hydrolyzed to the title compound in 50 % yield after chromatography.

5

Example 66

4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoic acid

10 Step 1: A solution of EtMgBr in ether (3 N, 2.17 eq) was cooled to - 70°C. The product of step 2 in Example 62 (1 eq) in ether (0.55 M) was added and the reaction mixture was stirred at - 70°C for 2 h. After the addition of benzoyl chloride (3 eq) in ether (1.5 M) the reaction was stirred at - 40°C for 2 h, quenched with saturated NaHCO₃ at - 40°C and allowed to warm to 25°C. EtOAc/water work up and crystallization from hexane/EtOAc the desired ketone in 89 % yield.

15

Step 2: The material from step 1 (1 eq) in CH₂Cl₂ (0.25 M) was treated with NEt₃ (2 eq), followed by BOC anhydride (1.24 eq) and DMAP (0.21 eq). After stirring for 20 minutes at 25°C the reaction mixture was worked up with CH₂Cl₂ and water. Pure desired material was obtained in 97 % yield by trituration with hexane.

20

Step 3: The above material (1 eq) in THF (0.3 M) was combined with pyridine (excess) and HFpyridine (excess) at 0°C. The reaction was stirred at 25°C for 1.5 h. EtOAc/water work up followed by chromatographic purification afforded the desired alcohol in 86 % yield.

25

Step 4: The alcohol from step 3 (1 eq) in CH₂Cl₂ (0.4 M) was treated with 2,6-lutidine (2.5 eq) followed by SOCl₂ (1.2 eq). After 30 minutes at 25°C the reaction was worked up with EtOAc and water. The crude product was purified by chromatography and trituration with hexane to afford the corresponding chloride in 30 75 % yield.

- 148 -

5 Step 5: A mixture of the above material (1 eq), 325 mesh K_2CO_3 (2.4 eq), b-naphthol (1.2 eq) and KI (1.2 eq) in CH_3CN (0.3 M) was heated to reflux for 2 h. EtOAc/water work up, followed by trituration in hexane/EtOAc and chromatography of the mother liquor yielded 70 % of the expected ether.

10 Step 6: A NaOMe solution in MeOH was prepared by dissolving Na (3 eq) in MeOH (0.2 M). The product of step 5 (1 eq) in THF (0.04 M) was added and the reaction mixture was stirred at 25°C for 3 h. EtOAc/water work up followed by trituration with hexane/EtOAc afforded the indole compound in 93 %.

15 Step 7: A solution of the product from step 6 (1 eq) in dry DMF (0.1 M) was treated with NaH (60 % in mineral oil, 1.1 eq). Methyl 4-bromobutyrate (1.2 eq) and KI (1.2 eq) were added after 1 h and the reaction mixture was stirred at 75°C for 3 h. EtOAc/water work up followed by chromatographic purification yielded 96 % of the required ester.

20

Step 8: The material from step 7 (1 eq) was hydrolyzed by the action of $LiOH \cdot H_2O$ (1.2 eq) in THF/MeOH/water (3/1/1, 0.2 M). After stirring at 25°C for 2 h, the reaction mixture was quenched with 1N HCl and extracted with EtOAc and CH_2Cl_2 . The organic extracts were combined, washed, dried and concentrated. The crude material was purified by trituration in hexane/EtOAc to afford the title compound in 86 % yield.

25

Example 67

30 3-[(4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)amino]benzoic acid

- 149 -

5 Step 1: The product of Example 66 (1 eq) was reacted with EDCI (1.37 eq) and methyl 3-amino benzoate (1.05 eq) in DMF (0.09 M) in the presence of DMAP (0.2 eq). The reaction was stirred at 25°C for 1.5 h. EtOAc/water work up, followed by flash chromatography produced the desired amide in 81 % yield.

10 Step 2: An analogous procedure to step 2 of Example 66 yielded 98 % of the title compound after purification by trituration in hexane/EtOAc.

Example 68

15 **4-[3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl]-N-[3-
{[(trifluoromethyl)sulfonyl]amino}carbonyl]phenyl]butanamide**

Step 1: Following step 1 of Example 67 and using the acid of Example 67 and the appropriate sulfonamide provided after overnight reaction and analogous work up the desired product. Trituration with hexane/EtOAc yielded 100 % of the title compound.

20

Example 69

**4-[4-[3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-
yl]butanoyl]amino]benzoic acid**

25 Step 1: Following step 1 of Example 67 above using acid of Example 66 and the appropriate aniline yielded 76 % of the expected amide.

Step 2: An analogous procedure to step 2 of Example 67 yielded 78 % of the title compound after purification by trituration in hexane/EtOAc.

30 **Example 70**

- 150 -

5 2-[(4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)amino]benzoic acid

Step 1: Following step 1 of Example 67 using acid of Example 66 and the appropriate aniline yielded 36 % of the expected amide after chromatography.

10 Step 2: An analogous procedure to step 2 of Example 67 yielded 67 % of the title compound after purification by trituration in hexane/EtOAc.

Example 71

15 3-[(4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)amino]propanoic acid

Step 1: Following step 1 of Example 67 using acid of Example 66 and the appropriate aniline yielded 96 % of the expected amide after chromatography.

20 Step 2: An analogous procedure to step 2 of Example 67 yielded 90 % of the title compound after purification by trituration in hexane/EtOAc.

Example 72

25 3-[(4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)amino]propanoic acid

Following step 1 of Example 67 using acid of Example 66 and the appropriate sulfonamide yielded 100 % of the title compound.

Example 73

30 N-(4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}butanoyl)-2-methylbenzenesulfonamide

- 5 Following step 1 of Example 67 using acid of Example 66 and the appropriate sulfonamide yielded 80 % of the title compound after chromatography.

Example 74

10 **5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}pentanoic acid**

- Step 1: A solution of the product from step 6 of Example 66 (1 eq) in dry DMF (0.1 M) was treated with NaH (60 % in mineral oil, 1.1 eq). Ethyl 4-bromopentanoate (1.2 eq) and KI (1.2 eq) were added after 1 h and the reaction mixture was stirred at
15 75°C for 2 h. EtOAc/water work up followed by chromatographic purification yielded 92 % of the required ester.

- Step 2: The material from step 1 (1 eq) was hydrolyzed by the action of LiOH·H₂O (1.3 eq) in THF/MeOH/water (3/1/1, 0.2 M). After stirring at 25°C for 3.5 h, the
20 reaction mixture was quenched with 1N HCl and extracted with EtOAc and CH₂Cl₂. The organic extracts were combined, washed, dried and concentrated. The crude material was purified by trituration in hexane/EtOAc to afford the title compound in 95 % yield.

25 **Example 75**

3-[(5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}pentanoyl)amino]benzoic acid

- Step 1: The acid of Example 74 (1 eq) was reacted with EDCI (1.37 eq) and methyl 3-amino benzoate (1.05 eq) in DMF (0.09 M) in the presence of DMAP (0.2 eq). The
30 reaction was stirred at 25°C for 2.5 h. EtOAc/water work up, followed by flash chromatography produced the desired amide in 78 % yield.

- 152 -

5

Step 2: An analogous procedure to step 2 of Example 74 yielded 83 % of the title compound after purification by trituration in hexane/EtOAc.

Example 76

10

**5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}-N-[3-
({[(trifluoromethyl)sulfonyl]amino}carbonyl)phenyl]pentanamide**

Step 1: Following step 1 of Example 68 using acid of Example 75 and the appropriate sulfonamide yielded 83 % of the title compound after overnight reaction and analogous work up.

15

Example 77

2-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}acetic acid

Step 1: Following step 1 of Example 74 using the appropriate bromide yielded 80 % of the expected amide after chromatography.

20

Step 2: An analogous procedure to step 2 of Example 74 yielded 90 % of the title compound after trituration in hexane/EtOAc.

Example 78

25

**(E)-4-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}-2-
butenoic acid**

Step 1: Following step 1 of Example 74 using the appropriate bromide yielded 33 % of the expected amide after chromatography.

30

Step 2: An analogous procedure to step 2 of Example 74 yielded 70% of the title compound after trituration in hexane/EtOAc.

- 153 -

5

Example 79

3-({3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl)methyl}benzoic acid

Step 1: Following step 1 of Example 74 using the appropriate bromide yielded 75 %
10 of the expected amide after chromatography.

Step 2: An analogous procedure to step 2 of Example 74 yielded 92 % of the title compound after trituration in hexane/EtOAc.

15

Example 80

1-{1-[4-(1,3-benzothiazol-2-ylcarbonyl)benzyl]-5-(benzylsulfanyl)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-3-yl}-1-ethanone

Step 1 The thioether from step 6, Example 62 was dissolved in methylene chloride (0.12 M), and under nitrogen at 0°C treated with methylmagnesium bromide (1.1 eq),
20 and stirred at 0 to 25°C for 1 h. The mixture was recooled to 0°C and acetyl chloride (1.17 eq) was added. After 1 h at 0°C, the reaction mixture was quenched by addition of half saturated ammonium chloride solution, and worked up with methylene chloride. Chromatographic purification afforded the pure product (37%).

25 Step 2 Following the procedure for Example 81 step 1 and using p-Toluoyl chloride yielded the corresponding para amide.

Step 3 Following the procedure for Example 81 step 2 and using the product from step 1 above, the desired product was obtained.

30

10076079-1021203
2021203-10076079

- 154 -

- 5 Step 4 Following the procedure for step 3, Example 81 and using the material from step 3 yielded the desired benzyl bromide.

Step 5 Following the procedure for Example 81 step 4, the material from step 4 was coupled to the material from step 1 to yield the titled compound.

10

Example 81

1-{1-[3-(1,3-benzothiazol-2-ylcarbonyl)benzyl]-5-(benzylsulfanyl)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-3-yl]-1-ethanone

- 15 Step 1 m-Toluoyl chloride (0.8 M) was added to triethylamine (2.44 eq) and methoxymethyl amine HCl (1.1 eq) dissolved in methylene chloride at 0°C over 20 min. The reaction was allowed to warm to 25°C. After stirring at 25°C for 1 day, workup with methylene chloride and water afforded crude product in ca. 100% yield.

- 20 Step 2 Under anhydrous conditions benzothiazole was dissolved in THF (0.35 M). At -78 °C added BuLi (1.1 eq). After 1 h at -78°C, added the amide from step 1 in THF, over 15 min. The reaction was allowed to warm to 25°C. After stirring at 25°C for 1 day, workup with ethyl acetate and water and chromatography afforded pure tolyl ketone product (52%).

- 25 Step 3 The tolyl ketone from step 2 was dissolved in carbon tetrachloride (0.19M), and NBS (1.2 eq) and AIBN (0.11 eq) were added. After 1 d at 60°C, about 1:1 of starting material and product were present. Resubmission under the same conditions, followed by filtration and recrystallization from ethyl acetate afforded pure bromobenzyl ketone product (28%).

30

- 155 -

- 5 Step 4 The indole from step 1 of Example 80 was dissolved in dry DMF (0.04 M), followed by NaH (1.25 eq). After 45 min at 25°C, added the bromobenzyl ketone from step 3 (1.25 eq), and stirred for 1 h at 25°C. Workup, chromatography, and recrystallization from ethyl acetate/hexanes afforded the pure title compound (45%).

10 **Example 82**

2-[3-({3-acetyl-5-(benzyloxy)-2-[(2-naphthylsulfanyl)methyl]-1H-indol-1-yl)methyl}benzoyl]-1,3-benzothiazole-6-carboxylic acid

- Step1 Ethyl 4-aminobenzoate was dissolved in 95% v/v HOAc/water (0.4 M) and at 25°C treated with sodium thiocyanate (4.1 eq), and stirred at 25°C for 20 min. The mixture was cooled to 5°C and bromine (1.17 eq) in 95% v/v HOAc/water was added. After 1 h at 0 °C, the reaction mixture was quenched by addition of water, and filtered. Workup (ethyl acetate and sodium bicarbonate solution) gave 81% of the pure thiocyanate product.

- 20 Step 2 The aryl thiocyanate from step 1 was treated with sodium sulfide nonahydrate dissolved in water (1.2 eq, 1.3 M), and heated at reflux for 45 min. The cooled mixture was filtered, acidified to pH = 6 with HOAc, extracted with ethyl acetate, and concentrated to give the thiophenol product (91%).

- 25 Step 3 The thiophenol from step 2 was dissolved in 90% v/v HCOOH/water (3.3 M), and zinc dust (cat.) was added. After 3 h at reflux, workup with ethyl acetate and alkali afforded pure benzothiazole ester (60%).

- Step 4 Benzothiazole ester (step 3) was dissolved in THF/methanol/water (8:3:3), (0.34 M), lithium hydroxide (2 eq) was added, and the mixture was stirred for 2 h at 25°C. Workup (ethyl acetate/ aqueous acid) afforded pure benzothiazole acid (100%).
- 30

- 156 -

5

Step 5 Under anhydrous conditions the benzothiazole acid (step 4) was dissolved in THF (0.052 M). At -78°C added BuLi (2.2 eq). After 1 h at -78°C to 0°C, added the amide (1.28 eq) from step 1 of Example 81, in THF, at -78°C, over 5 min. After 0.5 h at -78°C, the reaction was allowed to warm to 25°C. After stirring at 25°C for 2h, workup with ethyl acetate and water and chromatography afforded pure tolyl ketone acid product (64%).

10

Step 6 The tolyl ketone acid from step 5 was suspended in (1:1) THF/ethanol, (0.075 M), and conc. sulfuric acid (excess) was added. After reflux for 1 d, workup (ethyl acetate and sodium bicarbonate solution) and purification by chromatography gave the pure tolyl ketone ester product (69%).

15

Step 7 The tolyl ketone ester from step 6 was dissolved in carbon tetrachloride (0.05M), and NBS (1.2 eq) and AIBN (0.15 eq) were added. After 1.25 h at reflux, another portion of NBS (0.3 eq) and AIBN (0.07 eq) were added. Filtration and recrystallization from ethyl acetate afforded pure bromobenzyl ketone product (22%).

20

Step 8 The indole from step 1 of Example 80 was dissolved in dry DMF (0.06 M), followed by NaH (1.11 eq). After 45 min at 25°C, added the bromobenzyl ketone from step 7 above (1.25 eq), and stirred for 1 h at 25°C. Workup afforded the crude ester, used in the next step.

25

Step 9 The crude ester from step 8 was dissolved in THF/methanol/water (8:2:2), (0.013 M), lithium hydroxide (4.3 eq) was added in portions, as the mixture was stirred for a total of 3-4 d at 25°C. Workup (ethyl acetate/ aqueous HCl) and purification by chromatography afforded pure benzothiazole acid (31%).

30

- 157 -

5

Example 83

5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}-2-oxopentanoic acid

10 Step 1 The indole from step 6 of Example 66 was dissolved in dry DMF (0.17 M), followed by NaH (1.2 eq). After 1.5 h at 25°C, added 3-iodo-1-chloropropane, and stirred for 4 h at 25°C. Workup (ethyl acetate/bicarbonate solution) and trituration (ethyl acetate/hexanes) afforded the product in 89% yield.

15 Step 2 The alkyl chloride from step 1 was dissolved in methyl ethyl ketone (0.036 M), followed by NaI (1.6 eq). After stirring for 1 d at reflux in the dark, workup (ethyl acetate/water) afforded the product iodide in 97% yield.

20 Step 3 NaH (1.0 eq) was weighed into a dry flask under nitrogen, and dry benzene (0.14 M) was added. At 0°C, added dry DMF (0.4 M), ethyl 2-carboxy-1,3-dithiane (1.0 eq), and the iodide from step 2 (1.0 eq) into a separate dry flask. This DMF solution was then added at 0°C to the benzene suspension, and the mixture was allowed to warm to 25°C and stirred for 3 h at 25°C. Workup (ethyl acetate/water) and chromatography afforded the product dithianyl ester in 54% yield.

25

Step 4 Silver nitrate (4.5 eq) and NCS (4.0 eq) were dissolved in 4:1 v/v acetonitrile/water (ca. 0.04 M), and a solution of the dithianyl ester from step 3 (1 eq) in acetonitrile (0.03 M) was added at 25°C. After stirring for 5 min at 25°C, added a sodium sulfite solution, followed in one minute by a sodium carbonate solution.
30 Workup (ethyl acetate/water) and chromatography afforded the product ketoester in 17% yield.

- 158 -

5

Step 5 The ketoester from step 4 was dissolved in THF/water (8:1), (0.03 M), lithium hydroxide (1.5 eq) was added, and the mixture was stirred for 2 h at 25°C. Workup (ethyl acetate/ aqueous HCl) and purification by chromatography afforded pure ketoacid (64%).

10

Example 84

3-[(5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl}-2-oxopentanoyl)amino]benzoic acid

15 Step 1 The dithianyl ester from step 3 Example 83 was dissolved in THF/ethanol/water (5:2:2) (0.008 M), and lithium hydroxide (11 eq) was added, and the mixture was stirred for 1 d at 50°C. Workup (ethyl acetate/ aqueous HCl) and chromatography afforded the pure dithianyl acid (90%).

20 Step 2 The dithianyl acid from step 1 was dissolved in dry methylene chloride (0.08 M), and DMF (cat.). Oxalyl chloride (1.2 eq) was added, and the reaction was stirred at 25°C for 0.5 h. Concentration was followed by redissolution in dry methylene chloride, and addition , at 0°C, of methyl 3-aminobenzoate (1.05 eq) and triethylamine (1.0 eq). The reaction was warmed to 25 °C, and stirred there for 3 h. Workup (ethyl acetate/aqueous acid) and purification by chromatography afforded the
25 product dithianyl ester in 89% yield.

30 Step 3 The dithianyl ester from step 2 was dissolved in THF/methanol/water (6:4:3) (0.02 M), and lithium hydroxide (4.3 eq) was added, and the mixture was stirred for 3 h at 25°C. Workup (ethyl acetate/ aqueous HCl) afforded the dithianyl acid (R = 3-COOH) (91%).

- 159 -

5 Step 4 Silver nitrate (4.6 eq) and NCS (4.0 eq) were dissolved in 4:1 v/v
acetonitrile/water (ca. 0.03 M), and a solution of the dithianyl acid from step 3 (1 eq)
in acetonitrile (0.009 M) was added at 25°C. After stirring for 10 min at 25°C, there
was added a sodium sulfite solution, followed in one minute by a sodium carbonate
solution. Workup (ethyl acetate/aqueous acid) and chromatography afforded the title
10 compound in 82% yield.

Example 85

4-[(5-{3-benzoyl-5-(benzyloxy)-2-[(2-naphthyloxy)methyl]-1H-indol-1-yl})-2-oxopentanoyl)amino]benzoic acid

15 Step 1 The dithianyl acid from step 1 Example 84 was treated as described above for
step 2 Example 84 except ethyl 4-aminobenzoate was substituted. Purification by
chromatography afforded the product dithianyl ester ($R = 4\text{-COOC}_2\text{H}_5$) in 30% yield.

20 Step 2 The dithianyl ester from step 1 was treated as described for step 3 of Example
84. Purification by chromatography afforded the product dithianyl acid ($R = 4\text{-COOH}$) in 89% yield.

25 Step 3 The dithianyl from step 6 was treated as described for step 4 of Example 84.
Repeated purification by chromatography followed by trituration in pentane afforded
the title compound in 30 % yield.

Example 86

30 Table I reports data for the compounds described in the examples above in
cPLA2 inhibition assays (described below). In the data columns of the tables, assay
results are reported as an "IC₅₀" value, which is the concentration of a compound
which inhibits 50% of the activity of the phospholipase enzyme in such assay. Where

- 160 -

5 no numerical IC₅₀ value appears, "NA" denotes that inhibitory activity was not detected from such compound in the corresponding assay and a blank box denotes that the compound was not tested in such assay as of the time of filing of the present application.

10 Activity Assays

(a) Vesicle Assay

1-palmitoyl-2-[¹⁴C] arachidonyl phosphatidylcholine (58 mCi/mmol) (final concentration 6 M) and 1,2-dioleoylglycerol (final concentration 3 M) were mixed and dried under a stream of nitrogen. To the lipids was added 50 mM Hepes pH 7.5 (2x final concentration of lipids) and the suspension was sonicated for 3 min. at 4°C. To the suspension was added 50 mM Hepes pH 7.5, 300 mM NaCl, 2 mM DTT, 2 mM CaCl₂ and 2 mg/ml bovine serum albumin (BSA) (Sigma A7511) (1.2x final concentration of lipids). A typical assay consisted of the lipid mixture (85 l) to which was added consecutively, the inhibitor (5 l in DMSO) and cPLA₂, 10 ng for an automated system or 1 ng for a manual assay, in 10l of the BSA buffer. This assay was conducted by either the manual assay or automated assay protocol described below.

(b) Soluble Substrate Assay (LysoPC)

25 1-[¹⁴C]-palmitoyl-2-hydroxyphosphatidyl-choline (57 mCi/mmol) (final concentration 4.4 M) was dried under a stream of nitrogen. The lipid was resuspended by vortexing 80 mM Hepes pH 7.5, 1 mM EDTA (1.2 x final concentration). A typical assay consisted of lipid suspension (85 l) to which was added consecutively the inhibitor (5l in DMSO) and cPLA₂, 200 ng in 80 mM Hepes pH 7.5, 2 mM DTT and 1 M EDTA. This assay was conducted by either the manual assay or automated assay protocol described below.

- 161 -

5

(c) Automated Assay

The lipid suspension and inhibitor were pre-incubated for 7 min. at 37°C. Enzyme was added and the incubation was continued for a further 30 mins. The reaction was then quenched by the addition of decane: isopropanol: trifluoroacetic acid (192:8:1 w/v, 150 l). A portion of the quench layer (50 l) was passed through a Rainin Spheric-5 silica column (5, 30 x 2.1 mm) eluting with heptane:methanol:TFA (97:3:0.1 v/v). The level of [¹⁴C]- arachidonic acid was analyzed by an in-line Radiomatic Flo-One/Beta counter (Packard).

15

(d) Manual Assay

The lipid, inhibitor and enzyme mixture were incubated at 37°C for 30 min. The reaction was quenched by the addition of heptane:isopropanol:0.5M sulfuric acid (105:20:1 v/v, 200 l). Half of the quench layer was applied to a disposable silica gel column (Whatman SIL, 1 ml) in a vacuum manifold positioned over a scintillation vial. Free [¹⁴C]- arachidonic acid was eluted by the addition of ethyl ether (1 ml). The level of radioactivity was measured by liquid scintillation counter.

(e) PMN Assay

PMNs were isolated using Ficoll-Hypaque according to the manufacturers directions. Red blood cells contaminating the PMNs were removed by hypotonic lysis, and the PMN pellet was washed once, and resuspended in Hanks buffered saline at a concentration of 2×10^6 cells/ml. The cells were preincubated with inhibitors for 15 min at 37°C and then stimulated with 2 uM A23187. When monitoring LTB₄ production as a measure of cPLA₂ inhibition, the reaction was quenched with an equal volume of ice cold phosphate buffered saline. Cells were removed by centrifugation, and the LTB₄ present in the cell supernatant was measured

- 162 -

5 using the LTB₄ scintillation proximity assay provided by Amersham according to the
manufacturers directions. In the assays reported in the Tables above, LTB₄ was
measured. When monitoring arachidonic acid production, the reaction was quenched
with methanol containing D8-arachidonic acid as an internal reference. The lipids
were extracted by the method of Bligh et al. ((1959) Can. J. Biochem. Physiol., 37,
10 911-917), and the fatty acid was converted to the pentafluorobenzyl ester and
analyzed by GC-MS in a manner similar to that reported by Ramesha and Taylor
((1991) *Anal. Biochem.* 192, 173-180).

(f) RBL Assay

15 RBL-2H3 cells were routinely cultured as 37°C in a 5% CO₂ atmosphere in
minimal essential medium containing nonessential amino acids and 12% fetal calf
serum. The day before the experiment, cells were seeded into spinner flasks at 3 x
10⁵ cells/ml and 100 ng/ml DNP specific-IgE was added. After 20 hrs, the cells were
harvested by centrifugation and washed once in serum-free minimal essential media,
20 and resuspended to 2 x 10⁶ cells/ml in serum free media. The cells were then
preincubated with either inhibitor in DMSO (1% v/v) or DMSO (1% v/v) for 15 min
at 37°C followed by stimulation with DNP-BSA (300 ng/ml). After 6 min, the cells
were removed by centrifugation, and the supernatant was assayed for PGD₂ content in
accordance with known methods.

25 (g) Coumarine Assay

7-hydroxycoumarinyl 6-heptenoate was used as a monomeric substrate for
cPLA2 as reported previously (Huang, Z. et al., 1994, *Nalytical Biochemistry* 222,
110-115). Inhibitors were mixed with 200 µL assay buffer (80 mM Heped, pH 7.5, 1
30 mM EDTA) containing 60 µM 7-hydroxycoumarinyl 6-heptenoate. The reaction was
initiated by adding 4 µg cPLA2 in 50 µL assay buffer. Hydrolysis of the 7-

- 163 -

5 hydroxycounarimyl 6-heptenoate ester was monitored in a fluorometer by exciting at 360 nm and monitoring emission at 460 nm. Enzyme activity is proportional to the increase in emission at 460 nm per minute. In the presence of a cPLA2 inhibitor, the rate of increase is less.

10 Example 87

Compounds of the present invention were also tested for *in vivo* activity in a rat paw edema test according to the procedure described below. The results are reported in Table II.

15 Rat Carrageenan-Induced Footpad Edema Test

Each compound was suspended in 0.3ml absolute ethanol, 0.1 ml Tween-80 and 2.0 ml Dulbecco's PBS (without calcium or magnesium). To this mixture, 0.1ml 1N NaOH was added. After solution was complete, additional amounts of PBS were added to adjust the concentration to 1 mg/ml. All compounds remained in solution.

20 Compounds were administered i.v. in a volume of 5 ml/kg to male Sprague Dawley rats at the same time that edema was induced by injection of 0.05ml of 1% Type IV carrageenan into the hind footpad. Footpad volume was measured before dosing with compound and 3 hours after dosing with carrageenan.

25 Table 1: Inhibition in the LysoPC Assay

Example	PERCENT INHIBITION @	CONCENTRATION (micromolar)
1	50	15
1	50	17
2	50	50
2	50	57
3	8	50

- 164 -

3	50	158
4	50	30
4	23	50
4	50	53
5	50	8
5	50	9.5
6	13	20
6	50	28
7	8	20
7	50	40
7	50	50
8	50	25
8	50	38
9	43	50
9	50	56
9	50	70
10	50	42
10	44	50
10	50	50
11	56	12.5
11	50	13
11	50	19
11	50	25
11	50	33
11	43	50
12	50	140
12	50	160
13	50	160
14	50	35
14	50	42
15	50	7
15	50	7.5
16	50	33
16	50	36
17	50	12.5
17	50	17
17	50	23
18	50	6
18	50	6.5

1007009 001000

- 165 -

18	50	6.5
18	50	7
18	50	8
19	50	18
19	50	20
20	50	6
20	50	6
20	50	6.25
20	50	7
21	53	6.25
21	50	13
21	89	25
22	50	25
23	50	3.8
23	50	3.8
23	50	4
23	50	5.5
23	50	4.5
23	50	3
23	50	11
23	50	12.2
24	50	6
24	57	6.25
24	50	4
25	0	0
25	23	400
26	50	7
26	50	8
26	50	9
26	50	9
26	44	10
26	50	10
26	50	14
27	50	14
27	50	17
28	50	6.2
28	50	15
29	70	2.5
29	50	5

10075079-021202

- 166 -

30	50	7
30	50	8
30	50	8
30	50	12.5
30	50	15
31	50	33
31	50	36
32	50	6
32	50	6
32	50	6.25
32	50	7
33	50	18
33	50	20
34	50	28
34	50	38
34	50	55
35	60	5
35	50	6.5
35	50	10
35	50	18
36	50	5
36	50	12
36	50	18
36	50	27
37	50	15
37	50	18
37	50	32
37	50	36
38	50	6
38	50	8
38	50	13
38	50	15
39	50	18
39	50	19
39	50	20
40	50	6
40	50	6.5
40	50	6.5
40	50	7

20021201 152052001

- 167 -

	40	50	8
	41	50	5
	41	50	7
	42	53	6.25
	42	50	13
	42	89	25
	43	50	6
	43	50	6.5
	44	50	8
	44	50	10
	44	50	20
	45	50	6
	45	50	7.8
	46	50	2.5
	46	50	2.8
	47	50	6
	47	50	7
	47	50	7.5
	48	50	5.8
	49	50	2.3
	49	50	2.4
	49	56	2.5
	50	-23	30
	50	7	200
	50	38	400
	51	0	100
	51	15	200
	51	22	400
	52	50	90
	52	50	99
	52	50	99
	53	50	175
	53	50	180
	53	50	180
53A		-19	20
53A		50	125
53A		50	165
53B		19	20
53B		50	31

1003609 004000

- 168 -

53B	50	52
54	14	100
54	50	200
54	50	200
55	14	50
55	50	150
56	8	50
56	50	235
57	50	18
57	50	19
58	50	45
58	50	48
59	64	6.2
59	50	13
59	50	13
59	50	16
59	50	17
60	50	12.5
61	50	15
62	64	6.25
62	50	3.8
62	50	2.5
63	67	6.25
63	50	15
63	50	3.5
64	50	42
64	50	43
64	50	40
65	50	18
65	50	19.5
66	50	12.5
66	50	16
67	50	4
67	70	6.25
68	50	4.2
68	80	6.25
69	50	11.5
69	50	12
69	50	16

10075079-001000

- 169 -

70	50	7
70	50	9
71	3	20
71	15	20
72	50	2
72	50	3.8
72	50	6
72	50	7.5
73	38	20
73	50	20
74	50	16
75	50	5.7
75	66	6.25
76	50	4.5
76	80	6.25
77	53	12.5
77	50	15
78	50	12
78	50	17
78	50	26
79	50	6.25
79	50	8.5
80	50	46
80	50	46
81	47	50
81	48	50
82	50	3.8
83	65	6.25
83	50	8
83	50	9
84	50	4.5
84	50	8
85	50	7
85	50	10

- 170 -

5 Table 2: Inhibition in the Rat CPE Model

Example	ROUTE of ADMIN.	DOSE (mg/Kg)	PERCENT INHIBITION
1	IV	5	10.47
1	IV	5	15.27
2	IV	5	15.99
2	IV	5	-0.46
3	IV	5	9.27
4	IV	5	5.3
5	IV	5	10.08
5	IV	5	2.67
6	IV	5	11.14
7	IV	5	13.72
8	IV	5	6.5
9	PO	2	10.77
9	PO	10	12.65
10	PO	2	4.51
10	PO	10	12.1
11	PO	2	30.89
11	PO	10	27.76
12	IV	5	14.77
13	IV	5	13.96
14	IV	5	17.02
15	IV	5	-0.48
15	IV	5	3.01
15	IV	5	6
17	IV	5	11.59
17	IV	5	-8.02
19	IV	5	7.43
20	IV	5	3.25
20	IV	5	18.88
20	IV	5	9.39
20	IV	5	30.06
21	IV	5	6.95
21	IV	5	15.26
26	IV	5	8

- 172 -

53B	53 IV	5	0.36
	IV	5	3.61
	54 IV	5	0.71
	55 IV	5	5.56
	56 IV	5	-1.37
	57 IV	5	6.18
	58 IV	5	10.37
	66 IV	5	6.88
	68 IV	5	-4.65
	68 IV	5	10.7
	69 IV	5	17.37
	70 IV	5	18.23
	71 IV	5	17.03
	72 IV	5	3.52
	72 IV	5	10.82
	76 IV	5	23.66
	76 IV	5	20.64
	76 IV	5	-8.83
	77 IV	5	10.46
	77 IV	5	11.98
	78 IV	5	-2.56
	79 IV	5	-2.09
	79 IV	5	15.98

5

The compounds of this invention inhibit Cytosolic Phospholipase A2 (cPLA2) activity which is required for supplying arachidonic acid substrate to cyclooxygenase -1or 2 and 5-lipoxygenase, which in turn initiates the production of prostaglandins and leukotrienes, respectively. In addition, cPLA2 activity is essential for producing the lyso-phospholipid precursor to Platelet Activating Factor (PAF). Thus, these compounds are useful in the treatment and prevention of disease states in which leukotrienes, prostaglandins or PAF are involved. Moreover, in diseases where more than one of these agents plays a role, a cPLA2 inhibitor is efficacious than

- 173 -

- 5 leukotriene, prostaglandin or PAF receptor antagonists and also more effective than cyclooxygenase or 5-lipoxygenase inhibitors.

Therefore, the compounds, pharmaceutical compositions and regimens of the present invention are useful in treating and preventing the disorders treated by
10 cyclooxygenase-2, cyclooxygenase-1, and 5-lipoxygenase inhibitors and also are antagonists of the receptors for PAF, leukotrienes or prostaglandins. Diseases treatable by compounds, formulations and regimens of this invention include, but are not limited to, pulmonary disorders including diseases such as asthma, chronic bronchitis, and related obstructive airway diseases; allergies and allergic reactions
15 such as allergic rhinitis, contact dermatitis, allergic conjunctivitis, and the like; inflammation such as arthritis or inflammatory bowel diseases; skin disorders such as psoriasis, atopic eczema, acne, ultraviolet (UV) damage, burns and dermatitis; cardiovascular disorders such as atherosclerosis, angina, myocardial ischaemia, hypertension, platelet aggregation, and the like; and renal insufficiency induced by
20 immunological or chemical. The drugs may also be cytoprotective, preventing damage to the gastrointestinal mucosa by noxious agents. The compounds are also useful in the treatment of adult respiratory distress syndrome, endotoxin shock and ischaemia induced injury including myocardial or brain injury.

- 25 These compounds are especially useful in the treatment of arthritic disorders, including but not limited to rheumatoid arthritis, spondyloarthropathies, gouty arthritis, osteoarthritis, systemic lupus erythematosus and juvenile arthritis. The compounds of this invention are further useful in the treatment of post-operative inflammation, including that following ophthalmic surgery such as cataract surgery or
30 refractive surgery

- 174 -

5

The compounds of this invention can be used as an antipyretic agent and in combination with other antipyretic agents known in the art.

10 The compounds of this invention may be utilized in methods of treating pain, particularly the pain associated with inflammation. Specific methods include, but are not limited to, those for treating centrally mediated pain, peripherally mediated pain, musculo-skeletal pain, lumbosacral pain, structural or soft tissue injury related pain, progressive disease related pain, such as oncology and degenerative disorders, neuropathic pain, which can include both acute pain, such as acute injury or trauma,
15 pre- and post-surgical, migraine pain, dental pain, etc., chronic pains, such as neuropathic pain conditions of diabetic peripheral neuropathy, post-herpetic neuralgia and fibromyalgia, and inflammatory conditions such as osteoarthritis or rheumatoid arthritis, sequela to acute injury or trauma and cancer-related pain.

20 Compositions and compounds of this invention are also useful in the treatment of menstrual cramps, preterm labor, tendonitis, bursitis, allergic neuritis, cytomegalovirus infection, apoptosis, including HIV- induced apoptosis, lumbago, liver disease including hepatitis.

25 The methods and compositions herein are also useful in treating gastrointestinal conditions such as inflammatory bowel disease, Crohn's disease, gastritis, irritable bowel syndrome and ulcerative colitis and for the prevention of treatment of cancer such as colorectal cancer. The compounds and compositions of the present invention are also useful for the prevention or treatment of benign and
30 malignant tumors/neoplasia including cancers such as colorectal cancer, brain cancer,

- 175 -

5 bone cancer, epithelial cell-derived neoplasia (epithelial carcinoma) such as basal cell carcinoma, adenocarcinoma, gastrointestinal cancer, including lip cancer, mouth cancer, esophageal cancer, small bowel cancer and stomach cancer, colon cancer, liver cancer, bladder cancer, pancreatic cancer, ovarian cancer, cervical cancer, lung cancer, breast cancer, and skin cancers, such as squamous cell and basal cell cancers,
10 prostate cancer, renal cell carcinoma, and other known cancers that effect epithelial cells throughout the body. Neoplasias for which compositions of the invention are particularly useful are gastrointestinal cancer, Barrett's esophagus, liver cancer, bladder cancer, pancreas cancer, ovarian cancer, prostatic cancer, cervical cancer, lung cancer, breast cancer, and skin cancer, such as squamous cell and basal cell
15 cancers. The compounds and methods of this invention can also be used to treat the fibrosis occurring with radiation therapy. Such compositions can be used to treat subjects having adenomatous polyps, including those with familial adenomatous polyposis (FAP). Additionally, such compositions can be used to prevent polyps from forming in patients at risk of FAP. Compounds of this invention will be useful in the
20 treatment of cancers based on anti-angiogenic effects.

Further uses of this invention include treating inflammation in such diseases as vascular diseases, migraine headaches, periarteritis nodosa, thyroiditis, aplastic anemia, Hodgkin's disease, sclerodoma, rheumatic fever, type I diabetes,
25 neuromuscular junction disease including myasthenia gravis, white matter disease including multiple sclerosis, sarcoidosis, nephrotic syndrome, Behcet's syndrome, polymyositis, gingivitis, nephritis, hypersensitivity, swelling occurring after injury including brain edema, myocardial ischemia, and the like. Also included are treatment of ophthalmic diseases, such as retinitis, conjunctivitis, retinopathies,
30 uveitis, ocular photophobia, and of acute injury to the eye tissue. Treatments herein

- 176 -

5 of pulmonary inflammation, such as that associated with viral infections and cystic
fibrosis, and in bone resorption such as that accompanying osteoporosis. These
compounds and compositions are useful for the treatment of certain central nervous
system disorders, such as cortical dementias including Alzheimer's disease,
neurodegeneration, and central nervous system damage resulting from stroke,
10 ischemia and trauma. The compounds of this invention may also be useful in the
treatment of Parkinson's disease.

It will be understood that methods of treating or preventing the maladies listed
herein comprise administering to a mammal subject to or experiencing the malady,
15 which may also be referred to as a mammal in need thereof, a pharmaceutically
effective amount of a compound of this invention, or a pharmaceutically acceptable
salt thereof.

Methods of treating pain comprise administering to a mammal subject to such
20 pain a pharmaceutically effective amount of a compound of this invention alone or in
combination with one or more additional pharmaceutically effective agents for the
treatment of pain or inflammation or the related underlying medical condition.
Examples of drug agents which may be combined with the present compounds are
analgesics, anti-angiogenic agents, anti-neoplastic agents, These compounds may also
25 be combined with anti-epileptic compounds that have pain alleviating properties, such
as gabapentin and pregabalin.

One such combination method of this invention comprises administering to a
mammal in need thereof a pharmaceutically effective amount of a compound of this

- 177 -

5 invention and a pharmaceutically effective amount of a nontoxic N-methyl-D-aspartate (NMDA) receptor antagonist and/or an agent that blocks at least one major intracellular consequence of NMDA receptor activation. Examples of NMDA receptor antagonists useful in these methods include dextromethorphan, dextrorphan, amantadine and memantine, or the pharmaceutically acceptable salts thereof.

10

Another method herein of treating inflammation and inflammatory disorders comprises the co-administration to a mammal in need thereof of an inhibitor of induced nitric oxide synthase with a compound of this invention. Administration of this combination is useful for prophylactic or therapeutic administration in a mammal
15 experiencing or subject to an abnormally low level of nitric oxide synthase (NOS) activity, particularly those subject to hypertension or an elevated risk of pulmonary hypertension, ischemic stroke, myocardial infarction, heart failure, progressive renal disease, thrombosis, reperfusion injury, or a nervous system degenerative disorder, such as Alzheimer's disease, or those chronically exposed to hypoxic conditions.

20

The methods of this invention also include those for treating or preventing a neoplasia disorder in a mammal, including a human, in need of such treatment or prevention. The method comprises treating the mammal with a therapeutically effective amount of a compound of this invention in combination with an MMP
25 inhibitor. These two components may further be optionally combined with one or more agents selected from an antiangiogenesis agent, an antineoplastic agent, an adjunctive agent, an immunotherapeutic agent, an analgesic agent; and/or a radiotherapeutic agent. One such multiple component therapy comprises

- 178 -

- 5 administering to the mammal in need thereof a compound of this invention, a matrix metalloproteinase inhibitor and an antineoplastic agent.

The methods and combinations of this invention may be used for the treatment or prevention of neoplasia disorders including acral lentiginous melanoma, actinic
10 keratoses, adenocarcinoma, adenoid cystic carcinoma, adenomas, adenosarcoma, adenosquamous carcinoma, astrocytic tumors, bartholin gland carcinoma, basal cell carcinoma, bronchial gland carcinomas, capillary, carcinoids, carcinoma, carcinosarcoma, cavernous, cholangiocarcinoma, chondrosarcoma, choroid plexus papilloma/carcinoma, clear cell carcinoma, cystadenoma, endodermal sinus tumor,
15 endometrial hyperplasia, endometrial stromal sarcoma, endometrioid adenocarcinoma, ependymal, epitheloid, Ewing's sarcoma, fibrolamellar, focal nodular hyperplasia, gastrinoma, germ cell tumors, glioblastoma, glucagonoma, hemangioblastomas, hemangioendothelioma, hemangiomas, hepatic adenoma, hepatic adenomatosis, hepatocellular carcinoma, insulinoma, intraepithelial neoplasia,
20 interepithelial. squamous cell neoplasia, invasive squamous cell carcinoma, large cell carcinoma, leiomyosarcoma, lentigo maligna melanomas, malignant melanoma, malignant mesothelial tumors, medulloblastoma, medulloepithelioma, melanoma, meningeal, mesothelial, metastatic carcinoma, mucoepidermoid carcinoma, neuroblastoma, neuroepithelial adenocarcinoma nodular melanoma, oat cell
25 carcinoma, oligodendroglial, osteosarcoma, pancreatic polypeptide, papillary serous adenocarcinoma, pineal cell, pituitary tumors, plasmacytoma, pseudosarcoma, pulmonary blastoma, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, sarcoma, serous carcinoma, small cell carcinoma, soft tissue carcinomas, somatostatin-secreting tumor, squamous carcinoma, squamous cell carcinoma,
30 submesothelial, superficial spreading melanoma, undifferentiated carcinoma, uveal

- 179 -

5 melanoma, verrucous carcinoma, vipoma, well differentiated carcinoma, and Wilm's tumor.

Antineoplastic agents useful in the combination therapies herein include anastrozole, calcium carbonate, capecitabine, carboplatin, cisplatin, Cell Pathways
10 CP-461, docetaxel, doxorubicin, etoposide, fluorouracil, fluoxymestrine, gemcitabine, goserelin, irinotecan, ketoconazole, letrozol, leucovorin, levamisole, megestrol, mitoxantrone, paclitaxel, raloxifene, retinoic acid, tamoxifen, thiotepa, topotecan, toremifene, vinorelbine, vinblastine, vincristine, selenium (selenomethionine), ursodeoxycholic acid, sulindac sulfone, exemestane and eflornithine (DFMO), 1-[4-
15 (2-Azepan-1-yl-ethoxy)-benzyl]-2-(4-hydroxy-phenyl)-3-methyl-1H-indol-5-ol (TSE-424) and 2-(4-Hydroxy-phenyl)-3-methyl-1-(4-(2-piperidin-1-yl-ethoxy)-benzyl)-1H-indol-5-ol (ERA-923).

This invention also includes methods of utilizing the compounds herein in
20 combination with a proteinaceous interleukin-1 inhibitor, such as an IL-1 receptor antagonist (IL-lra), for preventing or treating inflammatory diseases in a mammal. Acute and chronic interleukin-1 (IL-1)-mediated inflammatory diseases of interest in these methods include, but is not limited to acute pancreatitis; ALS; Alzheimer's disease; cachexia/anorexia; asthma; atherosclerosis; chronic fatigue syndrome, fever;
25 diabetes (e.g., insulin diabetes); glomerulonephritis; graft versus host rejection; hemohorrhagic shock; hyperalgesia, inflammatory bowel disease; inflammatory conditions of a joint, including osteoarthritis, psoriatic arthritis and rheumatoid arthritis; ischemic injury, including cerebral ischemia (e.g., brain injury as a result of trauma, epilepsy, hemorrhage or stroke, each of which may lead to

- 180 -

5 neurodegeneration); lung diseases (e.g., ARDS); multiple myeloma; multiple
sclerosis; myelogenous (e.g., AML and CML) and other leukemias; myopathies (e.g.,
muscle protein metabolism, esp. in sepsis); osteoporosis; Parkinson's disease; pain;
pre-term labor; psoriasis; reperfusion injury; septic shock; side effects from radiation
therapy, temporal mandibular joint disease, tumor metastasis; or an inflammatory
10 condition resulting from strain, sprain, cartilage damage, trauma, orthopedic surgery,
infection or other disease processes.

202120 62052001
15 This invention also provides a method of administering one or more of the
compounds of this invention to a female in need thereof to substantially prevent or
reducing changes in the female's reproductive system associated with onset or
continuation of labor. Also provided is a method of substantially preventing or
reducing uterine contractility either occurring during pregnancy or associated with
menorrhagia. These methods may optionally include coadministration of a compound
of this invention with a progestogen, a progestin or a progestational agent.

20 All patent and literature references cited herein are incorporated as if fully set
forth herein.